

Introducing a new approach for comparing fuzzy quantities by Rank and Mode

R. Saneifard ^{* †}, M. Sabouri [‡]

Abstract

Many methods for ranking fuzzy numbers have been proposed. The existing methods for ranking generalized fuzzy numbers based on the optimistic index (α) gives different values for comparing the numbers by using different values of the optimistic index (α) and it is the shortcoming of this method. So we introduce a new defuzzification using a crisp number for ordering and comparing the fuzzy numbers. The calculation of the proposed method is simpler and easier in comparison the other methods and it provides the correct ordering of generalized fuzzy numbers. It is shown that the proposed modification satisfies all the reasonable properties of fuzzy quantities (I) and (II) proposed by X. Wang, E. E. Kerre [18, 19].

Keywords : Fuzzy numbers; Fuzzy function; L-R type generalized fuzzy number.

1 Introduction

IN decision analysis under the fuzzy environment ranking fuzzy numbers is a very important decision-making procedure. L-R fuzzy number as the most general form of fuzzy number has been used extensively. The purpose of this paper is to introduce a new method for ranking fuzzy numbers. Ranking fuzzy numbers were first proposed by Jain for decision making in fuzzy situations by representing the ill-defined quantity as a fuzzy set. Ranking fuzzy numbers is important in decision making, data analysis, artificial intelligence, economic systems and operations research. Jain [11, 12], Dubios and Prade [13] introduced the relevant concepts of fuzzy numbers. Bortlan and Degani [14] reviewed some methods to rank fuzzy numbers, Chen and Hwang [15] proposed fuzzy multiple attribute decision making, Choobineh and Li [16] proposed an index for ordering fuzzy numbers. The rest of this paper is organized as follows. Section 2 introduces basic concepts and definitions of fuzzy numbers. Section 3 proposes an approach to ranking fuzzy numbers based on RM method. Section 4

presents some numerical examples to illustrate the advantages of the proposed approach.

2 Preliminaries

In this section some basic definitions and arithmetic operations are reviewed.

Definition 2.1 *The characteristic function μ_A of a crisp set $A \subseteq X$ assigns a value either 0 or 1 to each member in X .*

This function can be generalized to a function μ_A such that the value assigned to the element of the universal set X fall within a specified range i.e. $\mu_A : X \rightarrow [0, 1]$. The assigned value indicate the membership grade of the element in the set A .

The function μ_A is called the membership function and the set $A = \{x \in X : \mu_A(x) > 0\}$ defined by μ_A for each $x \in X$ is called a fuzzy set [1, 2, 3, 4, 5, 6].

Definition 2.2 *An extended fuzzy number A is described as any fuzzy subset of the universe set U with membership function μ_A defined as follows [7, 8, 9, 10, 21]:*

- (a) μ_A is a continuous mapping from U to the closed interval $[0, w]$, $0 < w < 1$.
- (b) $\mu_A = 0$, for all $x \in (-\infty, a_1]$.

*Corresponding author. srsaneifard@yahoo.com

[†]Department of Mathematics, Urmia Branch, Islamic Azad University, Oroumieh, Iran.

[‡]Department of Mathematics, Urmia Branch, Islamic Azad University, Oroumieh, Iran.

- (c) μ_A is strictly increasing on $[a_1, a_2]$.
- (d) $\mu_A(x) = w$, for all $x \in [a_2, a_3]$ as w is constant and $0 < w \leq 1$.
- (e) μ_A is strictly decreasing on $[a_3, a_4]$.
- (f) $\mu_A(x) = 0$, for all $x \in [a_4, \infty)$.

Definition 2.3 A triangular fuzzy number A is a fuzzy number with a piecewise linear membership function μ_A defined by:

$$\mu_A = \begin{cases} \frac{x-a_1}{a_2-a_1}, & a_1 \leq x \leq a_2, \\ \frac{a_2-x}{a_3-a_2}, & a_2 \leq x \leq a_3, \\ 0, & \text{otherwise.} \end{cases} \quad (2.1)$$

Which can be denoted as a triplet (a_1, a_2, a_3) [14].

Definition 2.4 A trapezoidal fuzzy number A is a fuzzy number with a membership function μ_A defined by:

$$\mu_A = \begin{cases} \frac{x-a_1}{a_2-a_1}, & a_1 \leq x \leq a_2, \\ 1, & a_2 \leq x \leq a_3, \\ \frac{a_2-x}{a_3-a_2}, & a_3 \leq x \leq a_4, \\ 0, & \text{otherwise.} \end{cases} \quad (2.2)$$

Which can be denoted as a quarter (a_1, a_2, a_3, a_4) [15].

Definition 2.5 A fuzzy number $A = (a, b, c, d; w)_{LR}$ is said to be a $L - R$ type generalized fuzzy number if its membership function is given by

$$\mu_A(x) = \begin{cases} wL(\frac{x-a}{b-a}), & a < x < b, \\ w, & b \leq x \leq c, \\ wR(\frac{x-c}{d-c}), & c < x < d, \end{cases} \quad (2.3)$$

where L and R are references functions [20].

Definition 2.6 A $L - R$ type generalized fuzzy number $A = (a, b, c, d; w)_{LR}$ is said to be a generalized trapezoidal fuzzy number if its membership function is given by

$$\mu_A(x) = \begin{cases} w(\frac{x-a}{b-a}), & a < x < b, \\ w, & b \leq x \leq c, \\ w(\frac{x-c}{d-c}), & c < x < d, \end{cases} \quad (2.4)$$

2.1 Arithmetic operations

In this subsection, arithmetic operations between two $L - R$ type generalized fuzzy numbers, defined on universal set of real numbers, are reviewed [20].

Let $A_1 = (a_1, b_1, c_1, d_1; w_1)_{LR}$, $A_2 = (a_2, b_2, c_2, d_2; w_2)_{LR}$, be two $L - R$ type generalized fuzzy numbers and $A_3 = (a_3, b_3, c_3, d_3; w_3)_{RL}$, be a $R - L$ type generalized fuzzy number then

- (i) $A_1 \oplus A_2 = (a_1 + a_2, b_1 + b_2, c_1 + c_2, d_1 + d_2; \min(w_1, w_2))$.
- (ii) $A_1 \ominus A_3 = (a_1 - d_3, b_1 - c_3, c_1 - b_3, d_1 - d_3; \min(w_1, w_3))$.
- (iii) $\lambda A_1 = \begin{cases} (\lambda a_1, \lambda b_1, \lambda c_1, \lambda d_1; w_1)_{LR}, & \lambda \geq 0, \\ (\lambda d_1, \lambda c_1, \lambda b_1, \lambda a_1; w_1)_{RL}, & \lambda < 0. \end{cases}$

3 Proposed approach

In this section, we will propose the ranking of fuzzy numbers associated with the RM method and the minimum value of a_1, a_2 .

An efficient approach for comparing the fuzzy numbers is by the use of ranking function [20], $\Re: F(R) \rightarrow R$, where $F(R)$ is a set of fuzzy numbers defined on set of real numbers, which maps each fuzzy number into the real line, where a natural order exists i.e.,

- (i) $A \succ B$ if and only if $\Re(A) > \Re(B)$,
- (ii) $A \prec B$ if and only if $\Re(A) < \Re(B)$,
- (iii) $A \sim B$ if and only if $\Re(A) = \Re(B)$.

Remark 3.1 For all fuzzy numbers A, B, C and D we have [18]

- (i) $A \succ B \implies A \oplus C \succ B \oplus C$,
- (ii) $A \prec B \implies A \ominus C \prec B \ominus C$,
- (iii) $A \sim B \implies A \oplus C \sim B \oplus C$,
- (iv) $A \succ B, C \succ D \implies A \oplus C \succ B \oplus D$.

3.1 Method to find values of $RM(A), RM(B)$ [20]

Let $A = (a_1, b_1, c_1, d_1; w_1)_{(L_1R_1)}$ and $B = (a_2, b_2, c_2, d_2; w_2)_{(L_2R_2)}$ be two $L-R$ type generalized fuzzy numbers then use the following steps to find the values of $RM(A)$ and $RM(B)$

Step 1. Find $w = \min(w_1, w_2)$.

Step 2. Let the membership function of A and B be $\mu_A(x)$ and $\mu_B(x)$, respectively, where,

$$\mu_A(x) = \begin{cases} 0, & -\infty < x \leq a_1, \\ w_1 L_1(\frac{b_1-x}{b_1-a_1}), & a_1 \leq x < b_1, \\ w_1, & b_1 \leq x \leq c_1, \\ w_1 R_1(\frac{x-c_1}{d_1-c_1}), & c_1 < x \leq d_1, \\ 0, & d_1 \leq x < \infty, \end{cases}$$

and

$$\mu_B(x) = \begin{cases} 0, & -\infty < x \leq a_2, \\ w_2 L_2(\frac{b_2-x}{b_2-a_2}), & a_2 \leq x < b_2, \\ w_2, & b_2 \leq x \leq c_2, \\ w_2 R_2(\frac{x-c_2}{d_2-c_2}), & c_2 < x \leq d_2, \\ 0, & d_2 \leq x < \infty, \end{cases}$$

where $\mu_A(x) \in [0, w_1]$ and $\mu_B(x) \in [0, w_2]$.

Let λ represent the membership value, which is common for both A and B , then $\lambda \in [0, w]$, where $w = \min(w_1, w_2)$.

Let $wL_1(\frac{b_1-x}{b_1-a_1}) = \lambda$ and

$$\begin{aligned} wR_1(\frac{x-c_1}{d_1-c_1}) &= \lambda \\ &\Rightarrow \frac{b_1-x}{b_1-a_1} \\ &= L_1^{-1}(\frac{\lambda}{w}) \\ &\Rightarrow x \\ &= b_1 - (b_1 - a_1)L_1^{-1}(\frac{\lambda}{w}) \end{aligned}$$

and

$$\begin{aligned} (\frac{x-c_1}{d_1-c_1}) &= R_1^{-1}(\frac{\lambda}{w}) \\ &\Rightarrow x \\ &= c_1 - (d_1 - c_1)R_1^{-1}(\frac{\lambda}{w}) \end{aligned}$$

and now

$$\begin{aligned} \Re(A) &= \alpha \int_0^w (b_1 - (b_1 - a_1)L_1^{-1}(\frac{\lambda}{w}))d\lambda \\ &+ (1-\alpha) \int_0^w (c_1 - (d_1 - c_1)R_1^{-1}(\frac{\lambda}{w}))d\lambda. \end{aligned}$$

Let

$$\frac{\lambda}{w} = h \Rightarrow d\lambda = wd h$$

$$\begin{aligned} \Re(A) &= w\alpha \int_0^1 (b_1 - (b_1 - a_1)L_1^{-1}(h))dh \\ &+ w(1-\alpha) \int_0^1 (c_1 - (d_1 - c_1)R_1^{-1}(h))dh. \end{aligned}$$

Similarity,

$$\begin{aligned} \Re(B) &= w\alpha \int_0^1 (b_2 - (b_2 - a_2)L_2^{-1}(h))dh \\ &+ w(1-\alpha) \int_0^1 (c_2 - (d_2 - c_2)R_2^{-1}(h))dh. \end{aligned}$$

Step 3. If $\Re(A) \neq \Re(B)$ then $RM(A) = \Re(A)$ and $RM(B) = \Re(B)$ otherwise

$$\begin{aligned} RM(A) &= mode(A) \\ &= \alpha \int_0^w b_1 dx + (1-\alpha) \int_0^w c_1 dx \\ &= w\alpha b_1 + (1-\alpha)w c_1, \end{aligned}$$

and

$$\begin{aligned} RM(B) &= mode(B) \\ &= \alpha \int_0^w b_2 dx + (1-\alpha) \int_0^w c_2 dx \\ &= w\alpha b_2 + (1-\alpha)w c_2, \end{aligned}$$

For generalized trapezoidal numbers $A = (a, b, c, d; w)$,

$$\Re(A) = \frac{w}{2}\alpha(a+b) + \frac{w}{2}(1-\alpha)(c+d),$$

and for triangular fuzzy numbers $A = (a, b, c; w)$,

$$\Re(A) = \frac{w}{2}\alpha(a+b) + \frac{w}{2}(1-\alpha)(b+c).$$

3.2 Method to find values of $RM_a(A)$ and $RM_a(B)$

In this subsection, we will propose a new method for ranking $L - R$ type generalized fuzzy numbers. So use the following steps to find the values of $RM_a(A)$ and $RM_a(B)$.

Step 1. Find $a = \min(a_1, a_2)$.

Step 2. Let $RM_a(A) = RM(A) - a$ and similarity,

Step 3. $RM_a(B) = RM(B) - a$.

(i) $A \succ B$ if and only if $RM_a(A) > RM_a(B)$.

(ii) $A \prec B$ if and only if $RM_a(A) < RM_a(B)$.

(iii) $A \sim B$ if and only if $RM_a(A) = RM_a(B)$.

We consider the following axioms for the ordering approach [18, 19].

A_1 : For an arbitrary finite subset Γ of E and $A \in \Gamma$, $A \succeq A$.

A_2 : For an arbitrary finite subset Γ of E and $(A, B) \in \Gamma^2$, $A \succeq B$ and $B \succeq A$, we should have $A \sim B$.

A_3 : For an arbitrary finite subset Γ of E and $(A, B, C) \in \Gamma^3$, $A \succeq B$ and $B \succeq C$, we should have $A \succeq C$.

\mathcal{A}_4 : For an arbitrary finite subset Γ of E and $(A, B) \in \Gamma^2$, $\inf \text{supp}(A) > \sup \text{supp}(B)$ we should have $A \succeq B$.

\mathcal{A}'_4 : For an arbitrary finite subset Γ of E and $(A, B) \in \Gamma^2$, $\inf \text{supp}(A) > \sup \text{supp}(B)$ we should have $A \succeq B$.

\mathcal{A}_5 : Let Γ and Γ' be two arbitrary finite subset of E in which A and B are in $\Gamma \cap \Gamma'$. We obtain the ranking order $A \succ B$ by defuzzification $RM_a(A)$ on Γ if only and if $A \succ B$ by $RM_a(A)$ on Γ' .

\mathcal{A}_6 : Let $A, B, A+B$ and $B+C$ be elements of E . If $A \succeq B$ then $A+C \succeq B+C$.

\mathcal{A}'_6 : Let $A, B, A+B$ and $B+C$ be elements of E . If $A \succeq B$ then $A+C \succ B+C$.

In addition for the axioms A_1, A_2, A_6 we can consider two following properties:

\mathcal{A}_7 : For an arbitrary finite subset Γ of E and $A \in \Gamma$, the defuzzification must belong to its support function.

\mathcal{A}_8 : For an arbitrary finite subset Γ of E and $A \in \Gamma$, distance between A and its defuzzification will be minimized.

Defuzzification

$$RM_a(A) : E \rightarrow R$$

is called the best approximation operator for any $A \in E$ if it has axioms A_1, A_2, \dots, A_8 .

4 Numerical example

In this section, some examples are used to illustrate the proposed approach to ranking $L-R$ fuzzy numbers.

Example 4.1 Let $A = (0.1, 0.3, 0.3, 0.5; 1)$ and $B = (0.2, 0.3, 0.3, 0.4; 1)$ be two generalized trapezoidal fuzzy numbers.

Find $a = \min(0.1, 0.2) = 0.1$.

For $\alpha = 0$, $RM(A) = 0.4$, $RM(B) = 0.35$, $RM_a(A) = 0.4 - 0.1 = 0.3$ and $RM_a(B) = 0.35 - 0.1 = 0.25$. So $A \succ B$.

For $\alpha = 1$, $RM(A) = 0.2$, $RM(B) = 0.25$, $RM_a(A) = 0.2 - 0.1 = 0.1$ and $RM_a(B) = 0.25 - 0.1 = 0.15$. So $A \prec B$.

For $\alpha = \frac{1}{2}$, $RM(A) = 0.3$, $RM(B) = 0.3$, $RM_a(A) = 0.3 - 0.1 = 0.2$ and $RM_a(B) = 0.3 - 0.1 = 0.2$. So $A \sim B$.

Example 4.2 Consider the three triangular fuzzy numbers $A = (6, 1, 1)$, $B = (6, 0.1, 1)$ and $C = (6, 0, 1)$. By using our method, $RM(A) = 1 + \frac{5}{2}\alpha$, $RM(B) = \frac{5}{2}\alpha + 0.55$, $RM(C) = \frac{5}{2}\alpha + \frac{1}{2}$ and with $a = \min(6, 6, 6) = 6$, we have $RM_a(A) = \frac{5}{2}\alpha - 5$, $RM_a(B) = \frac{5}{2}\alpha - 5.45$, $RM_a(C) = \frac{5}{2}\alpha - 5.5$. So $A \prec B \prec C$.

Example 4.3 Let $A = (0.2, 0.4, 0.6, 0.8; 0.35)$ and $B = (0.1, 0.2, 0.3, 0.4; 0.7)$ be two generalized trapezoidal fuzzy numbers.

For $a = \min(0.2, 0.1) = 0.1$, and $\alpha = 0$, then $RM(A) = 0.245$, $RM(B) = 0.1225$, $RM_a(A) = 0.245 - 0.1 = 0.145$ and $RM_a(B) = 0.1225 - 0.1 = 0.0225$. So $A \succ B$.

Also, for $\alpha = 1$ we have, $RM(A) = 0.105$, $RM(B) = 0.0525$, $RM_a(A) = 0.105 - 0.1 = 0.005$ and $RM_a(B) = 0.0525 - 0.1 = -0.0475$. So $A \succ B$.

For $\alpha = \frac{1}{2}$, $RM(A) = 0.175$, $RM(B) = 0.0875$, $RM_a(A) = 0.175 - 0.1 = 0.075$ and $RM_a(B) = 0.0875 - 0.1 = -0.0125$. So $A \succ B$.

5 Conclusion

This paper presents a new approach for ranking $L-R$ fuzzy numbers. The examples given in this paper illustrate that the proposed approach gives the correct ordering of fuzzy numbers. Comparing with the existing approaches, it is efficient and simple. For the validation of the proposed ranking function, it is shown that this ranking function satisfies all the reasonable properties of fuzzy quantities (I) and (II) proposed by X.Wang, E.E. Kerre [18, 19].

References

- [1] S. Abbasbandy, B. Asady, *Ranking of fuzzy numbers by sign distance*, Inform. Sci. 176 (2006) 2405-2416.
- [2] S. Abbasbandy, T. Hajjari, *A new approach for ranking of trapezoidal fuzzy numbers*, Computer and Mathematics with Appl. 57 (2009) 413-419.
- [3] R. Saneifard, T. Allahviranloo, F. Hosseinzadeh, N. Mikaeilvand, *Euclidean ranking DMUs with fuzzy data in DEA*, Applied Mathematical Sciences 60 (2007) 2989-2998.
- [4] R. Saneifard, *A method for defuzzification by weighted distance*, Int. J. Industrial Mathematics 3 (2009) 209-217.
- [5] R. Saneifard, *Ranking L-R fuzzy numbers with weighted averaging based on levels*, Int. J. Industrial Mathematics 2 (2009) 163-173.
- [6] R. Saneifard, *Some properties of neural networks in designing fuzzy systems*, Neural Comput. Appl. <http://dx.doi.org/10.1007/s00521-011-0777-1>.
- [7] R. Saneifard, *Designing an algorithm for evaluating decision-making units based on neural weighted function*, Neural Comput. Appl. <http://dx.doi.org/10.1007/s00521-012-0878-5>.

- [8] Rahim Saneifard and Rasoul Saneifard, *A modified method for defuzzification by probability density function*, Journal of Applied Sciences Research 2 (2011) 102-110.
- [9] Rahim Saneifard and Rasoul Saneifard, *Correlation Coefficient Between Fuzzy Numbers Based On Central Interval*, Journal of Fuzzy Set Valued Analysis <http://dx.doi.org/10.5899/2012/jfsva-00108/>.
- [10] Rahim Saneifard and Rasoul Saneifard, *The Median Value of Fuzzy Numbers and its Applications in Decision Making*, Journal of Fuzzy Set Valued Analysis <http://dx.doi.org/10.5899/2012/jfsva-00051/>.
- [11] R. Jain, *Decision-making in the presence of fuzzy variable*, IEEE Trans. Syst. Man Cybern. 6 (1976) 698-703.
- [12] R. Jain, *A procedure for multi-aspect decision making fuzzy sets*, Int. J. Syst. Sci. 8 (1997) 1-7.
- [13] D. Dubois, H. Prade, *Operations on fuzzy numbers*, International Journal of Systems Sciences 9 (1978) 613-626.
- [14] G. Bortlan, R. Degani, *A review of some methods for ranking fuzzy numbers*. Fuzzy Sets and Systems 15 (1985) 1-19.
- [15] S. J. Chen, C. L. Hwang, *Fuzzy multiple attribute decision making*, Springer, New York 1992.
- [16] F. Choobineh, H. Li, *An index for ordering fuzzy numbers*, Fuzzy Sets and Systems 54 (1993) 287-294.
- [17] G. Dias, *Ranking alternatives using fuzzy numbers: A computational approach*, Fuzzy Sets and Systems 56 (1993) 247-252.
- [18] X. Wang, E. E. Kerre, *Reasonable properties for the ordering of fuzzy quantities (I)*, Fuzzy Sets and Syst. 118 (2001) 375-385.
- [19] X. Wang, E. E. Kerre, *Reasonable properties for the ordering of fuzzy quantities (II)*, Fuzzy Sets and Syst. 118 (2001) 387-405.
- [20] A. Kumar, P. Singh, A. Kaur, P. Kaur, *RM approach for ranking of L-R type generalized fuzzy numbers*, Springer, Verlag 15 (2011) 1373-1381.
- [21] A. Kaufmann, M. M. Gupta, *Fuzzy mathematical models in engineering and management science*. Elsevier. Amsterdam 1988.



Rahim Saneifard was born in 1972 in Oroumieh, Iran. He received B.Sc (1997) in pure mathematics and M.Sc. in applied mathematics from Azarbi-jan Teacher Education University to Tabriz, Islamic Azad University Lahijan Branch to Lahijan, respectively. He is a Assistant Prof. in the department of mathematics at Islamic Azad University, Urmia Branch, Oroumieh, in Iran. His current interest is in fuzzy mathematics.



Mahsa Sabouri was born in 1983 in Oroumieh, Islamic Republic of Iran. She received the B.S. degree in applied mathematics from payam nor university of Oroumieh Branch and M.S. degree in applied mathematics from Islamic Azad University of Oroumieh in 2012.