

# A method for solving first order fuzzy differential equation

O. Sedaghatfar <sup>\* †</sup>, P. Darabi <sup>‡</sup>, S. Moloudzadeh <sup>§</sup>

---

## Abstract

In this paper, a new approach for solving first-order fuzzy differential equation (*FDE*) with fuzzy initial value under strongly generalized H-differentiability is considered. The idea of the presented approach is constructed based on the extending 0-cut and 1-cut solution of original *FDE*. First, under H-differentiability the solutions of fuzzy differential equations in 0-cut and 1-cut cases are found and convex combination of them, is considered. Then we choose the initial interval that 0-cut of original problem is positive. The obtained convex combination on this interval is the solution of *FDE*.

*Keywords* : Fuzzy differential equation (FDE); Interval differential equation; Strongly generalized H-differentiability.

---

## 1 Introduction

THE topic of fuzzy differential equation (*FDE*) has been rapidly growing in recent years. Kandel and Byatt [19] applied the concept of (*FDE*) to the analysis of fuzzy dynamical problems. The (*FDE*) and the initial value problem (Cauchy problem) were rigorously treated by Kaleva [17, 18], Seikkala [23], He and Yi [13], Kloeden [20] and by other researchers (see [3, 9, 10, 12, 16]). The numerical methods for solving *FDE* are introduced in [1, 2, 4, 5]. Bede [7] applied the concept of Strongly generalized H-differentiability to solving first order linear fuzzy differential equations and Allahviranloo et. al. [6] proposed a method to obtain analytical solutions for *FDE* under strongly generalized H-differentiability.

The idea of the presented approach is constructed based on the extending 0-cut and 1-cut solution of original *FDE*. Obviously, 0-cut of *FDE* is interval differential equation or ordinary differential equation. First, under H-differentiability *FDE* has been divided in two differential equations and solutions of each fuzzy differential equations in 0-cut and 1-cut cases are found. Then in each cases of differentiability, the initial interval that 0-cut of original problem is positive, is found. The obtained convex combination on this interval, is an solution of *FDE*.

The structure of this paper is organized as follows. In Section 2, some basic definitions and notations which will be used are brought. In Section 3, first order fuzzy differential equation is introduced and the proposed approach is given in detail. In Section 4, the proposed method is illustrated by solving several examples. Conclusion is drawn in Section 5.

---

\*Corresponding author. sedaghtfar@yahoo.com.com

<sup>†</sup>Department of Mathematics, Shahr-e-rey Branch, Islamic Azad University, Shahr-e-rey, Iran.

<sup>‡</sup>Department of Mathematics, Science and Research Branch, Islamic Azad University, Tehran, Iran.

<sup>§</sup>Department of Mathematics, Science and Research Branch, Islamic Azad University, Tehran, Iran.

## 2 Basic Definitions and Notations

**Definition 2.1** An arbitrary fuzzy number is represented by an ordered pair of functions  $(\underline{u}(r), \bar{u}(r))$  for all  $r \in [0, 1]$ , which satisfy the following requirements [15].

- $\underline{u}(r)$  is a bounded left continuous nondecreasing function over  $[0, 1]$ ;
- $\bar{u}(r)$  is a bounded left continuous non-increasing function over  $[0, 1]$ ;
- $\underline{u}(r) \leq \bar{u}(r)$ ,  $0 \leq r \leq 1$ .

Let  $E$  be the set of all upper semi-continuous normal convex fuzzy numbers with bounded  $r$ -level intervals. This means that if  $\tilde{v} \in E$  then the  $r$ -level set

$$[v]_r = \{s | v(s) \geq r\},$$

is a closed bounded interval which is denoted by  $[v]_r = [\underline{v}(r), \bar{v}(r)]$  for  $r \in (0, 1]$  and  $[v]_0 = \bigcup_{r \in (0, 1]} [v]_r$ .

Two fuzzy numbers  $\tilde{u}$  and  $\tilde{v}$  are called equal  $\tilde{u} = \tilde{v}$ , if  $u(s) = v(s)$  for all  $s \in \mathbb{R}$  or  $[u]_r = [v]_r$  for all  $r \in [0, 1]$ .

**Lemma 2.1** [22] If  $\tilde{u}, \tilde{v} \in E$ , then for  $r \in (0, 1]$ ,

$$[u + v]_r = [\underline{u}(r) + \underline{v}(r), \bar{u}(r) + \bar{v}(r)],$$

$$[u.v]_r = [\min k_r, \max k_r],$$

where

$$k_r = \{\underline{u}(r)\underline{v}(r), \underline{u}(r)\bar{v}(r), \bar{u}(r)\underline{v}(r), \bar{u}(r)\bar{v}(r)\}.$$

The Hausdorff distance between fuzzy numbers given by  $D : E \times E \rightarrow R^+ \cup \{0\}$ ,

$$D(u, v) = \sup_{r \in [0, 1]} \max\{|\underline{u}(r) - \underline{v}(r)|, |\bar{u}(r) - \bar{v}(r)|\},$$

where  $u = (\underline{u}(r), \bar{u}(r))$ ,  $v = (\underline{v}(r), \bar{v}(r)) \subset E$  is utilized (see [8]). Then it is easy to see that  $D$  is a metric in  $E$  where for all  $u, v, w, e \in E$  has the following properties (see [21]).

- $D(u \oplus w, v \oplus w) = D(u, v)$ ,
- $D(k \odot u, k \odot v) = |k|D(u, v)$ ,  $\forall k \in R$ ,
- $D(u \oplus v, w \oplus e) \leq D(u, w) + D(v, e)$ ,
- $(D, E)$  is a complete metric space.

**Definition 2.2** [14]. Let  $f : R \rightarrow E$  be a fuzzy valued function. If for arbitrary fixed  $t_0 \in R$  and  $\varepsilon > 0$ ,  $\delta > 0$  such that

$$|t - t_0| < \delta \implies D(f(t), f(t_0)) < \varepsilon,$$

$f$  is said to be continuous.

**Definition 2.3** Let  $x, y \in E$ . If there exists  $z \in E$  such that  $x = y + z$ , then  $z$  is called the  $H$ -difference of  $x$  and  $y$  and it is denoted by  $x \ominus y$ . In this paper we consider the following definition of differentiability for fuzzy-valued functions which was introduced by Bede et.al. [8] and investigate by Chalco-Cano et.al. [11].

**Definition 2.4** Let  $f : (a, b) \rightarrow E$  and  $x_0 \in (a, b)$ . We say that  $f$  is strongly generalized  $H$ -differentiable at  $x_0$ . If there exists an element  $f'(x_0) \in E$ , such that:

(1) for all  $h > 0$  sufficiently near to 0,  $\exists f(x_0 + h) \ominus f(x_0)$ ,  $\exists f(x_0) \ominus f(x_0 - h)$  such that the following limits hold.

$$\lim_{h \rightarrow 0^+} \frac{f(x_0+h) \ominus f(x_0)}{h} =$$

$$\lim_{h \rightarrow 0^+} \frac{f(x_0) \ominus f(x_0-h)}{h} = f'(x_0)$$

(2) for all  $h < 0$  sufficiently near to 0,  $\exists f(x_0) \ominus f(x_0 + h)$ ,  $\exists f(x_0 - h) \ominus f(x_0)$  such that the following limits hold.

$$\lim_{h \rightarrow 0^+} \frac{f(x_0) \ominus f(x_0+h)}{h} =$$

$$\lim_{h \rightarrow 0^+} \frac{f(x_0-h) \ominus f(x_0)}{h} = f'(x_0)$$

In the special case when  $f$  is a fuzzy-valued function, we have the following results.

**Theorem 2.1** [11]. Let  $f : R \rightarrow E$  be a function and denote  $f(t) = (\underline{f}(t; r), \bar{f}(t; r))$ , for each  $r \in [0, 1]$ . Then

- (1) if  $f$  is differentiable in the first form (1) in definition 2.4 then  $\underline{f}(t; r)$  and  $\bar{f}(t; r)$  are differentiable functions and  $f'(t) = (\underline{f}'(t; r), \bar{f}'(t; r))$ .
- (2) if  $f$  is differentiable in the second form (2) in definition 2.4 then  $\underline{f}(t; r)$  and  $\bar{f}(t; r)$  are differentiable functions and  $f'(t) = (\bar{f}'(t; r), \underline{f}'(t; r))$ .

The principal properties of the  $H$ -derivatives in the first form (1), some of which still hold for the second form (2), are well known and can be found in [17] and some properties for the second form (2) can be found in [11].

Notice that we say fuzzy-valued function  $f$  is (I)-differentiable if satisfies in the first form (1) in Definition 2.4 and we say  $f$  is (II)-differentiable if satisfies in the second form (2) in Definition 2.4.

### 3 First Order Fuzzy Differential Equations

In this section, we are going to investigate solution of *FDE*. Consider the following first order fuzzy differential equation:

$$\begin{cases} y'(t) = f(t, y(t)) \\ \tilde{y}(t_0) = \tilde{y}_0, \end{cases} \quad (3.1)$$

where  $f : [a, b] \times E \rightarrow E$  is fuzzy-valued function,  $\tilde{y}_0 \in E$  and strongly generalized H-differentiability is also considered which is defined in Definition 2.4

Now, we describe our propose approach for solving *FDE* (3.1). First, we shall solve *FDE* (3.1) in sense of 1-cut and 0-cut as a follows:

$$\begin{cases} (y')^{[1]}(t) = f^{[1]}(t, y(t)), \\ y^{[1]}(t_0) = \tilde{y}_0^{[1]}, \end{cases} \quad t_0 \in [0, T]. \quad (3.2)$$

$$\begin{cases} (y')^{[0]}(t) = f^{[0]}(t, y(t)), \\ y^{[0]}(t_0) = \tilde{y}_0^{[0]}, \end{cases} \quad t_0 \in [0, T]. \quad (3.3)$$

If Eq. (3.2) and Eq. (3.3) be a crisp differential equation we can solve it as usual, otherwise, if Eq. (3.2) and Eq. (3.3) be an interval differential equation we will solve it by stefanini et. al.'s method is proposed and discussed in [24]. Notice that the solutions of differential equation (3.2) and Eq. (3.3) are presented with notation  $y^{[1]}(t)$  and  $y^{[0]}(t)$  respectively. Then unknown  $\tilde{y}(t)$  must be determined, this approach leads to obtain

$$y(t) = [y(t; r), \bar{y}(t; r)] = [(1 - r)y^{[0]}(t) + ry^{[1]}(t), (1 - r)\bar{y}^{[0]}(t) + r\bar{y}^{[1]}(t)] \quad (3.4)$$

Pleased notice that, we assumed the 0-cut and 1-cut solutions are differentiable. Consequently, based on type of differentiability we have two following cases:

**Case I.** Suppose that  $\tilde{y}(t)$  in Eq. (3.3) is (I)-differentiable, then we get:

$$y'(t) = [y'(t; r), \bar{y}'(t; r)] \quad (3.5)$$

Consider Eq. (3.5) and original *FDE* (3.1), then we have the following for all  $r \in [0, 1]$ :

$$\begin{cases} \underline{y}'(t; r) = \underline{f}(t; r), & t_0 \leq t \leq T, \\ \bar{y}'(t; r) = \bar{f}(t; r), & t_0 \leq t \leq T. \end{cases} \quad (3.6)$$

Therefore

$$\begin{cases} (1 - r)(\underline{y}^{[0]})'(t) + r(\underline{y}^{[1]})'(t) = (1 - r)(\underline{f}^{[0]})(t) + r(\underline{f}^{[1]})(t), \\ (1 - r)(\bar{y}^{[0]})'(t) + r(\bar{y}^{[1]})'(t) = (1 - r)(\bar{f}^{[0]})(t) + r(\bar{f}^{[1]})(t), \\ \underline{y}(t_0; r) = (1 - r)\underline{y}^{[0]}(t_0) + r\underline{y}^{[1]}(t_0), \\ \bar{y}(t_0; r) = (1 - r)\bar{y}^{[0]}(t_0) + r\bar{y}^{[1]}(t_0). \end{cases} \quad (3.7)$$

Then we have

$$\begin{cases} \underline{y}^{[0]}'(t) = \underline{f}^{[0]}(t), \\ \bar{y}^{[0]}'(t) = \bar{f}^{[0]}(t), \\ \underline{y}^{[0]}(t_0) = \underline{y}_0^{[0]}, \\ \bar{y}^{[0]}(t_0) = \bar{y}_0^{[0]}, \end{cases} \quad (3.8)$$

and

$$\begin{cases} \underline{y}^{[1]}'(t) = \underline{f}^{[1]}(t), \\ \bar{y}^{[1]}'(t) = \bar{f}^{[1]}(t), \\ \underline{y}^{[1]}(t_0) = \underline{y}_0^{[1]}, \\ \bar{y}^{[1]}(t_0) = \bar{y}_0^{[1]}. \end{cases} \quad (3.9)$$

Indeed, we will find  $\underline{y}^{[0]}(t)$ ,  $\bar{y}^{[0]}(t)$ ,  $\underline{y}^{[1]}(t)$ ,  $\bar{y}^{[1]}(t)$  by solving *ODEs* (3.8), (3.9). Hence, solution of original *FDE* (3.1) is derived based on 1-cut solution and 0-cut solution as follows:

$$\tilde{y}(t) = [y(t; r), \bar{y}(t; r)] = [(1 - r)\underline{y}^{[0]}(t) + r\underline{y}^{[1]}(t), (1 - r)\bar{y}^{[0]}(t) + r\bar{y}^{[1]}(t)],$$

where for all  $0 \leq r \leq 1$  and  $t \in [0, T]$  such that:

$$\begin{aligned} \underline{y}(t; r) &= (1 - r)\underline{y}^{[0]}(t) + r\underline{y}^{[1]}(t), \\ \bar{y}(t; r) &= (1 - r)\bar{y}^{[0]}(t) + r\bar{y}^{[1]}(t). \end{aligned}$$

**Case II.** Suppose that  $\tilde{y}(t)$  in Eq. (3.3) is (II)-differentiable, then we get:

$$y'(t) = [\bar{y}'(t; r), \underline{y}'(t; r)] \quad (3.10)$$

Similarly, *ODEs* (3.8) and (3.9) can be rewritten in sense of (II)-differentiability as following:

$$\begin{cases} \underline{y}^{[0]'}(t) = \overline{f^{[0]}}(t), \\ \overline{y}^{[0]'}(t) = \underline{f^{[0]}}(t), \\ \underline{y}^{[0]}(t_0) = \overline{y_0^{[0]}}, \\ \overline{y}^{[0]}(t_0) = \underline{y_0^{[0]}}, \end{cases} \quad (3.11)$$

and

$$\begin{cases} \underline{y}^{[1]'}(t) = \overline{f^{[1]}}(t), \\ \overline{y}^{[1]'}(t) = \underline{f^{[1]}}(t), \\ \underline{y}^{[1]}(t_0) = \overline{y_0^{[1]}}, \\ \overline{y}^{[1]}(t_0) = \underline{y_0^{[1]}}. \end{cases} \quad (3.12)$$

Finally, by solving above *ODEs* (3.11) and (3.12)  $\underline{y}^{[0]}(t)$ ,  $\overline{y}^{[0]}(t)$ ,  $\underline{y}^{[1]}(t)$ ,  $\overline{y}^{[1]}(t)$  are determined and follows we can drive solution of original *FDE* (3.1) in sense of (II)-differentiability by using

$$\begin{aligned} \underline{y}(t; r) &= (1 - r)\underline{y}^{[0]}(t) + ry^{[1]}(t), \\ \overline{y}(t; r) &= (1 - r)\overline{y}^{[0]}(t) + r\overline{y}^{[1]}(t), \end{aligned}$$

for all  $0 \leq r \leq 1$  and  $t \in [0, T]$ .

### 4 Examples

In this section, some examples are given to illustrate our method and we show that our approach is coincide with the exact solutions.

**Example 4.1** Let consider the following *FDE*:

$$\begin{cases} y'(t) = y(t) + \tilde{a}, \\ y(0; r) = [r, 2 - r], \quad \tilde{a} = [r - 1, 1 - r] \quad 0 \leq r \leq 1. \end{cases} \quad (4.13)$$

**Case I.** Suppose that  $\tilde{y}(t)$  is (I)-differentiable. The exact solution of above system is:

$$\tilde{y}(t) = [(2r - 1)e^t - r + 1, -(2r - 3)e^t r - 1].$$

Based on *ODEs* (3.8) and (3.9), we have:

$$\begin{cases} \underline{y}^{[0]'}(t) = \underline{y}^{[0]}(t) - 1, \\ \overline{y}^{[0]'}(t) = \overline{y}^{[0]}(t) + 1, \\ \underline{y}^{[0]}(0) = 0, \\ \overline{y}^{[0]}(0) = 2, \end{cases} \quad (4.14)$$

$$\begin{cases} \underline{y}^{[1]'}(t) = \underline{y}^{[1]}(t), \\ \overline{y}^{[1]'}(t) = \overline{y}^{[1]}(t), \\ \underline{y}^{[1]}(0) = 1, \\ \overline{y}^{[1]}(0) = 1. \end{cases} \quad (4.15)$$

By solving *ODEs* (4.16) and (4.17), we get:

$$\begin{aligned} \underline{y}^{[0]}(t) &= -e^t + 1, \quad \overline{y}^{[0]}(t) = 3e^t - 1, \\ \underline{y}^{[1]}(t) &= e^t, \quad \overline{y}^{[1]}(t) = e^t. \end{aligned}$$

Finally, with substituting above solution in (3.4) we have:

$$\underline{y}(t) = (1 - r)(-e^t + 1) + re^t = e^t(2r - 1) - r + 1,$$

$$\overline{y}(t) = (1 - r)(3e^t - 1) + re^t = r - e^t(2r - 3) - 1,$$

and

$$\tilde{y}(t) = [e^t(2r - 1) - r + 1, r - e^t(2r - 3) - 1],$$

where  $y(t)$  has valid level sets for  $t > 0$  and  $y(t)$  is (I)-differentiable on  $t > 0$ , then  $y(t)$  is a solution for original problem on  $t > 0$ .

**Case II.** Suppose that  $\tilde{y}(t)$  is (II)-differentiable. The exact solution of above system is:

$$\tilde{y}(t) = [e^t - r + (2r - 2)/e^t + 1, r + e^t - (2r - 2)/e^t - 1].$$

Based on *ODEs* (3.8) and (3.9), we have:

$$\begin{cases} \underline{y}^{[0]'}(t) = \overline{y}^{[0]}(t) + 1, \\ \overline{y}^{[0]'}(t) = \underline{y}^{[0]}(t) - 1, \\ \underline{y}^{[0]}(0) = 0, \\ \overline{y}^{[0]}(0) = 2, \end{cases} \quad (4.16)$$

$$\begin{cases} \underline{y}^{[1]'}(t) = \overline{y}^{[1]}(t), \\ \overline{y}^{[1]'}(t) = \underline{y}^{[1]}(t), \\ \underline{y}^{[1]}(0) = 1, \\ \overline{y}^{[1]}(0) = 1. \end{cases} \quad (4.17)$$

By solving ODEs (4.16) and (4.17), we get:

$$\begin{aligned} \underline{y}^{[0]}(t) &= e^t - r + (2r - 2)/e^t + 1, \\ \overline{y}^{[0]}(t) &= r + e^t - (2r - 2)/e^t - 1, \\ \underline{y}^{[1]}(t) &= e^t, \quad \overline{y}^{[1]}(t) = e^t. \end{aligned}$$

Finally, with substituting above solution in (3.4) we have:

$$\underline{y}(t) = (1 - r)(e^t - r + (2r - 2)/e^t + 1) + re^t - (1/2)e^{-t} = e^t - r + (2r - 2)/e^t + 1,$$

$$\overline{y}(t) = (1 - r)(r + e^t - (2r - 2)/e^t - 1) + re^t = r + e^t - (2r - 2)/e^t - 1$$

and

$$\tilde{y}(t) = [e^t - r + (2r - 2)/e^t + 1, r + e^t - (2r - 2)/e^t - 1].$$

where  $y(t)$  has valid level sets for  $[0, \log(2)]$  and  $y(t)$  is (II)-differentiable on  $t > 0$ , then  $y(t)$  is a solution for the original problem on  $[0, \log(2)]$ .

**Example 4.2** Let consider the following FDE:

$$\begin{cases} y'(t) = -y(t), \\ y(0; r) = [1 + r, 5 - r], \quad 0 \leq r \leq 1. \end{cases} \quad (4.18)$$

**Case I.** Suppose that  $\tilde{y}(t)$  is (I)-differentiable. The exact solution of above system is:

$$\tilde{y}(t) = [3e^{-t} + (r - 2)e^t, 3e^{-t} - (r - 2)e^t].$$

Based on ODEs (3.11) and (3.12), we get:

$$\begin{cases} \underline{y}^{[0]'}(t) = -\overline{y}^{[0]}(t), \\ \overline{y}^{[0]'}(t) = -\underline{y}^{[0]}(t), \\ \underline{y}^{[0]}(0) = 1, \\ \overline{y}^{[0]}(0) = 5, \end{cases} \quad (4.19)$$

$$\begin{cases} \underline{y}^{[1]'}(t) = -\overline{y}^{[1]}(t), \\ \overline{y}^{[1]'}(t) = -\underline{y}^{[1]}(t), \\ \underline{y}^{[1]}(0) = 2, \\ \overline{y}^{[1]}(0) = 4. \end{cases} \quad (4.20)$$

By solving ODEs (4.21) and (4.22), we get:

$$\begin{aligned} \underline{y}^{[0]}(t) &= 3e^{-t} - 2e^t, \quad \overline{y}^{[0]}(t) = 3e^{-t} + 2e^t, \\ \underline{y}^{[1]}(t) &= 3e^{-t} - e^t, \quad \overline{y}^{[1]}(t) = 3e^{-t} + e^t. \end{aligned}$$

Finally, with substituting above solution in (3.4) we have:

$$\underline{y}(t) = (1 - r)(3e^{-t} - 2e^t) + r(3e^{-t} - e^t) = 3e^{-t} + (r - 2)e^t,$$

$$\overline{y}(t) = (1 - r)(3e^{-t} + 2e^t) + r(3e^{-t} + e^t) = 3e^{-t} - (r - 2)e^t$$

and

$$\tilde{y}(t) = [3e^{-t} + (r - 2)e^t, 3e^{-t} - (r - 2)e^t],$$

where  $y(t)$  has valid level sets for  $t > 0$  and  $y(t)$  is (I)-differentiable on  $t > 0$ , then  $y(t)$  is a solution for the original problem on  $t > 0$ .

**Case II.** Suppose that  $\tilde{y}(t)$  is (II)-differentiable. The exact solution of above system is:

$$\tilde{y}(t) = [(1 + r)e^{-t}, (5 - r)e^{-t}].$$

Based on ODEs (3.11) and (3.12), we get:

$$\begin{cases} \underline{y}^{[0]'}(t) = -\underline{y}^{[0]}(t), \\ \overline{y}^{[0]'}(t) = -\overline{y}^{[0]}(t), \\ \underline{y}^{[0]}(0) = 1, \\ \overline{y}^{[0]}(0) = 5, \end{cases} \quad (4.21)$$

$$\begin{cases} \underline{y}^{[1]'}(t) = -\underline{y}^{[1]}(t), \\ \overline{y}^{[1]'}(t) = -\overline{y}^{[1]}(t), \\ \underline{y}^{[1]}(0) = 2, \\ \overline{y}^{[1]}(0) = 4. \end{cases} \quad (4.22)$$

By solving ODEs (4.21) and (4.22), we get:

$$\begin{aligned} \underline{y}^{[0]}(t) &= e^{-t}, \quad \overline{y}^{[0]}(t) = 5e^{-t}, \\ \underline{y}^{[1]}(t) &= 2e^{-t}, \quad \overline{y}^{[1]}(t) = 4e^{-t}. \end{aligned}$$

Finally, with substituting above solution in (3.4) we have:

$$\underline{y}(t) = (1 - r)e^{-t} + 2re^{-t} = (1 + r)e^{-t},$$

$$\bar{y}(t) = 5(1 - r)e^{-t} + 4re^{-t} = (5 - r)e^{-t}$$

and

$$\tilde{y}(t) = [(1 + r)e^{-t}, (5 - r)e^{-t}],$$

where  $y(t)$  has valid level sets for  $t > 0$  and  $y(t)$  is (II)-differentiable on  $t > 0$ , then  $y(t)$  is a solution for original problem on  $t > 0$ .

## 5 Conclusion

A new method for solving first order fuzzy differential equations *FDE* with fuzzy initial value under strongly generalized H-differentiability was considered. The idea of the presented approach was constructed based on the extending 0-cut and 1-cut solution of original *FDE*. First, under H-differentiability the solutions of fuzzy differential equations in 0-cut and 1-cut cases were found then convex combination of them was considered. Then we found out the initial interval that 0-cut of original problem was positive. The obtained convex combination on this interval was a solution of *FDE*. Using 0-cut and 1-cut solutions we show that the discussed method can be applied to solve the fuzzy differential equation.

## References

- [1] S. Abbasbandy, T. Allahviranloo, *Numerical solutions of fuzzy differential equations by Taylor method*, Comput. Methods Appl. Math. 2 (2002) 113-124.
- [2] S. Abbasbandy, T. Allahviranloo, Óscar López-Pouso, J. J. Nieto, *Numerical Methods for Fuzzy Differential Inclusions*, Comput. Math. Appl. 48 (2004) 1633-1641.
- [3] T. Allahviranloo, S. Salahshour, *A new approach for solving first order fuzzy differential equations*, IPMU. (2010) 522-531.
- [4] T. Allahviranloo, N. Ahmady, E. Ahmady, *Numerical solution of fuzzy differential equations by predictorcorrector method*, Infor. Sci. 177 (2007) 1633-1647.
- [5] T. Allahviranloo, N. Ahmady, E. Ahmady, *A method for solving n-th order fuzzy linear differential equations*, Comput. Math. Appl. 86 (2009) 730-742.
- [6] T. Allahviranloo, N. A. Kiani, M. Barkhordari, *Toward the existence and uniqueness of solution of second-order fuzzy differential equations*, Information Sciences 179 (2009) 1207-1215.
- [7] B. Bede, I. J. Rudas, A. L. Bencsik, *First order linear fuzzy differential equations under generalized differentiability*, Inform. Sci. 177 (2007) 1648-1662.
- [8] B. Bede, S. G. Gal, *Generalizations of the differentiability of fuzzy-number-valued functions with applications to fuzzy differential equation*, Fuzzy Sets and Systems 151 (2005) 581-599.
- [9] J. J. Buckley, T. Feuring, *Fuzzy differential equations*, Fuzzy sets and Systems 110 (2000) 43-54.
- [10] J. J. Buckley, T. Feuring, *Introduction to fuzzy partial differential equations*, Fuzzy Sets and Systems 105 (1999) 241-248.
- [11] Y. Chalco-cano, H. Roman-Flores, *On new solutions of fuzzy differential equations*, Chaos, Solitons and Fractals 38 (2006) 112-119.
- [12] W. Congxin, S. Shiji, *Existence theorem to the Cauchy problem of fuzzy differential equations under compactness-type conditions*, Inf. Sci. 108 (1998) 123-134.
- [13] O. He, W. Yi, *On fuzzy differential equations*, Fuzzy Sets and Systems 24 (1989) 321-325.
- [14] M. Friedman, M. Ming, A. Kandel, *Numerical solution of fuzzy differential and integral equations*, Fuzzy Sets and System 106 (1999) 35-48.
- [15] M. Friedman, M. Ming, A. Kandel, *Fuzzy linear systems*, Fuzzy Set and Systems 96 (1998) 201-209.
- [16] L. J. Jowers, J. J. Buckley, K. D. Reilly, *Simulating continuous fuzzy systems*, Inf. Sci. 177 (2007) 436-448.



- [17] O. Kaleva, *Fuzzy differential equations*, Fuzzy Sets and Systems 24 (1987) 301-317.
- [18] O. Kaleva, *The Cauchy problem for fuzzy differential equations*, Fuzzy Sets and Systems 35 (1990) 389-396.
- [19] A. Kandel, W. J. Byatt, *Fuzzy differential equations*, In: Proceedings of the International Conference on Cybernetics and Society, Tokyo (1978) 1213-12160.
- [20] P. Kloeden, *Remarks on peano-like theorems for fuzzy differential equations*, Fuzzy Sets and Systems 44 (1991) 161-164.
- [21] M. L. Puri, D. Ralescu, *Fuzzy random variables*, J. Math. Anal. Appl. 114 (1986) 409-422.
- [22] D. Ralescu, *A survey of the representation of fuzzy concepts and its applications*, In: M. M. Gupta, R. K. Ragade, R. R. Yager, Eds., *Advances in Fuzzy Set Theory and Applications* (North-Holland, Amsterdam (1979) 77-91.
- [23] S. Seikkala, *On the fuzzy initial value problem*, Fuzzy Sets and Systems 24 (1987) 319-330.
- [24] L. Stefanini, B. Bede, *Generalized Hukuhara differentiability of interval-valued functions and interval differential equations*, Nonlinear Anal. 71 (2009) 1311-1328.



Omolbanin Sedaghatfar was born in the Tehran, Shahr-e-rey in 1982. She received B.Sc and M.Sc degree in applied mathematics from Zahedan Branch, Tehran - Science and Research Branch, *IAU*, respectively, and his Ph.D degree in applied mathematics from science and research Branch, *IAU*, Tehran, Iran in 2011. She is a Assistant Prof. in the department of mathematics at Islamic Azad University, Shahr-e-rey, Tehran in Iran. She is research interests include numerical solution of fuzzy integro-differential equations and fuzzy differential equation and similar topics.



Pejman Darabi was born in Tehran-Iran in 1978. He received B.Sc and M.Sc degrees in applied mathematics from Sabzevar Teacher Education University, science and research Branch, *IAU* to Tehran, respectively, and his Ph.D degree in applied mathematics from science and research Branch, *IAU*, Tehran, Iran in 2011. Main research interest include numerical solution of Fuzzy differential equation systems, ordinary differential equation, partial differential equation, fuzzy linear system and similar topics.



Saeid Moloudzadeh was born in the Naghadeh-Iran in 1976. He received B.Sc degree in mathematics and M.Sc degree in applied mathematics from Payam-e-Noor University of Naghadeh, science and research Branch, *IAU* to Tehran, respectively, and his Ph.D degree in applied mathematics from Science and Research Branch, *IAU*, Tehran, Iran in 2011. He is research interests include fuzzy mathematics, especially, on numerical solution of fuzzy linear systems, fuzzy differential equations and similar topics.