

Numerical solution of two-dimensional fuzzy Fredholm integral equations using collocation fuzzy wavelet like operator

N. Hassasi ^{*}, R. Ezzati ^{†‡}

Abstract

In this paper, first we propose a new method to approximate the solution of two-dimensional linear fuzzy Fredholm integral equations of the second kind based on the fuzzy wavelet like operator. Then, we discuss and investigate the convergence and error analysis of the proposed method. Finally, to show the accuracy of the proposed method, we present two numerical examples.

Keywords : Fuzzy linear system; Two-dimensional fuzzy Fredholm integral equation; Fuzzy wavelet like operator.

1 Introduction

THE concept of fuzzy integrals was initiated by Dubois and Prade [11] and then investigated by Kaleva [21], Goetschel and Voxman [20], Nanda [25] and others. In [35], the Henstock integral of fuzzy-valued functions is defined, while the fuzzy Riemann integral and its numerical integration was investigated by Wu in [36]. In [17], the authors introduced some quadrature rules for the integral of fuzzy-number-valued mappings. Kaleva [21] proposed the existence and uniqueness of the solution of fuzzy differential equations using the Banach fixed point principle. Mordeson and Newman [24], started the study of the subject of fuzzy integral equations (FIE).

Many authors applied the Banach fixed point principle, as a powerful tool, to show the existence and uniqueness of the solution of FIE [5, 6, 16, 29, 30, 31]. In [19, 26], sufficient conditions

for bounded solutions of FIE are given. Recently, the iterative techniques are applied to solve fuzzy Fredholm integral equations of the second kind (FFIE-2) by researchers [7, 15, 28]. Friedman et al. [16] presented a numerical algorithm to solve FFIE-2 based on successive approximations method. Also, Friedman et al. [17] investigated numerical procedures for solving such equations by using the embedding method. In [8], the successive approximations method is used for solving nonlinear fuzzy Fredholm integral equations. The authors of [44] presented iterative method and quadrature rules for solving nonlinear FFIE-2. In [9], Bica et al. developed an iterative numerical method to solve fuzzy Hammerstein-Volterra integral equations with constant delay. In [10], the same method has been applied to obtain the solution that take values in the set of right-sided fuzzy numbers for a fuzzy Volterra integral equation with constant delay arising in epidemiology. For numeric-analytic methods to solve FFIE-2, one can refer [1, 4, 12, 13, 14, 18, 27, 32, 33, 34, 43]. Since many real-valued problems in engineering and mechanics can be brought in the form of two-dimensional fuzzy Fredholm integral equa-

^{*}Department of Mathematics, Karaj Branch, Islamic Azad University, Karaj, Iran.

[†]Corresponding author. ezati@kiau.ac.ir

[‡]Department of Mathematics, Karaj Branch, Islamic Azad University, Karaj, Iran.

tions (2DFFIE), it is important that we develop quadrature rules and numerical methods for solving such equations. Recently, some researchers investigated solving such equations. In [?], the authors applied modified Homotopy perturbation method to solve 2DFFIE. The authors of [38] proposed quadratur rules for numerical solution of two-dimensional fuzzy integrals. In this work, the authors also obtained numerical solution of linear 2DFFIE by using iterative technique. Also, solving nonlinear 2DFFIE by using quadrature rules is done in [42].

Recently several researchers have attempted to develop "fuzzy wavelets" based models and systems. Wavelet theory is a relatively new and an emerging area in mathematical research. Also, wavelets are the suitable and powerful tool for approximating functions based on wavelet basis functions. As we know, most of the methods to solve integral equations lead us to solve the linear systems, but the singularity of these systems may cause some problems. So, by using fuzzy wavelet like operator with collocation points to obtain numerical solution of 2FFIE, such linear systems can be as sparse linear systems and it can reduce the cost of computation. In this paper, by using fuzzy wavelet like operator, we propose a numerical method to approximate the solution of linear 2FFIE.

$$f(x, y) = g(x, y) \oplus \lambda \int_c^d \int_a^b K(x, y, s, t) \otimes f(s, t) ds dt \tag{1.1}$$

where $K(x, y, s, t)$ is an arbitrary kernel function over the square $[a, b] \times [c, d]$, and $g(x, y)$ is a fuzzy real valued function of x and y . Also, we present the error estimation for approximating the solution of equation(1.1).

The rest of the paper is organized as follows: In Section 2, we review some elementary concepts of the fuzzy set theory and modulus of continuity. In Section 5, we present the method for approximate the solution of (1.1) by using fuzzy wavelet like operator. The error estimation of this method is proved in Section 4. Finally in Section 5, we give two numerical examples for applicability of the proposed method and compare the numerical results with the exact solutions.

2 Preliminaries

In this Section, we review some necessary backgrounds and notions of fuzzy sets theory.

Definition 2.1 [3] *A fuzzy number is a function $u : R \rightarrow [0, 1]$ with the following properties:*

1. u is normal, that is $\exists x_0 \in R$ such that $u(x_0) = 1$,
2. u is fuzzy convex set (i.e. $u(\lambda x + (1 - \lambda)y) \geq \min\{u(x), u(y)\} \quad \forall x, y \in R, \lambda \in [0, 1]$),
3. u is upper semicontinuous on R ,
4. The $\overline{\{x \in R : u(x) > 0\}}$ is compact set.

The set of all fuzzy numbers denoted by R_F .

Definition 2.2 [17] *Suppose that $u \in R_F$. The r -level set of u is denoted by $[u]^r = [u_-^{(r)}, u_+^{(r)}]$ and is defined by $[u]^r = \{x \in R; u(x) \geq r\}$, where $0 < r \leq 1$. Also, $[u]^0$ is called the support of u and it is given as $[u]^0 = \overline{\{x \in R : u(x) > 0\}}$. It follows that the r -level sets of u are closed and bounded intervals in R .*

It is well-known that the addition and multiplication operations of real numbers can be extended to R_F . In other words, for $u, v \in R_F$ and $\lambda \in R$, we define uniquely the sum $u \oplus v$ and the product $\lambda \otimes u$, by

$$[u \oplus v]^r = [u]^r + [v]^r, \quad [\lambda \otimes u]^r = \lambda [u]^r, \quad \forall r \in [0, 1]$$

where $[u]^r + [v]^r$ means the usual addition of two intervals (as subset of R) and $\lambda [u]^r$ means the usual product between a scalar and a subset of R . We use the same symbol \sum both for the sum of real numbers and for the sum \oplus (when the terms are fuzzy numbers).

Definition 2.3 [17] *An arbitrary fuzzy number is represented, in parametric form, by an ordered pair of functions $(\underline{u}(r), \bar{u}(r)), 0 \leq r \leq 1$, which satisfy the following requirements:*

1. $\underline{u}(r)$ is a bounded left continuous nondecreasing function over $[0, 1]$,
2. $\bar{u}(r)$ is a bounded left continuous nonincreasing function over $[0, 1]$,
3. $\underline{u}(r) \leq \bar{u}(r), \quad 0 \leq r \leq 1$.
4. The addition and scalar multiplication of fuzzy numbers in R_F are defined as follows:

$$u \oplus v = (\underline{u}(r) + \underline{v}(r), \bar{u}(r) + \bar{v}(r))$$

$$\lambda \otimes u = \begin{cases} (\lambda \underline{u}(r), \lambda \bar{u}(r)) & \lambda \geq 0 \\ \lambda \bar{u}(r), \lambda \underline{u}(r) & \lambda < 0. \end{cases}$$

Definition 2.4 [3] *For arbitrary fuzzy numbers $u = (\underline{u}, \bar{u})$ and $v = (\underline{v}, \bar{v})$ the quantity*

$$D(u, v) = \sup_{r \in [0, 1]} \max\{|u_-^{(r)} - v_-^{(r)}|, |u_+^{(r)} - v_+^{(r)}|\}$$

is called the distance between u and v . It is shown that (R_F, D) is a complete metric space with the following properties [7]:

1. $D(u \oplus w, v \oplus w) = D(u, v), \forall u, v, w \in R_F,$
2. $D(k \otimes u, k \otimes v) = |k|D(u, v), \forall u, v \in R_F, \forall k \in R,$
3. $D(u \oplus v, w \oplus e) \leq D(u, w) + D(v, e), \forall u, v, e \in R_F.$

Definition 2.5 [11] Let $f, g : [a, b] \times [c, d] \rightarrow R_F$ be fuzzy number valued functions. The uniform distance between f and g is defined by

$$D^*(f, g) = \sup\{D(f(x, y), g(x, y)); x \in [a, b], y \in [c, d]\}.$$

Definition 2.6 [40] Let $f, g : [a, b] \times [c, d] \rightarrow R_F$. For each partition $p = \{x_1, x_2, \dots, x_n\}$ of $[a, b]$ and $q = \{y_1, y_2, \dots, y_n\}$ of $[c, d]$ and for arbitrary $\xi_i : x_{i-1} \leq \xi_i \leq x_i, 2 \leq i \leq n$ and for arbitrary $\eta_j : y_{j-1} \leq \eta_j \leq y_j, 2 \leq j \leq n$, let

$$R_p = \sum_{i=2}^m \sum_{j=2}^n f(\xi_i, \eta_j) \otimes (x_i - x_{i-1})(y_j - y_{j-1}).$$

The function f is called two-dimensional Riemann integrable to $I \in R_F$ if for every $\epsilon > 0$,

$$D(I, R_p) < \epsilon$$

In this case, we have

$$I = (FR) \int_c^d \int_a^b f(x, y) dx dy.$$

Definition 2.7 [42] A function $f : [a, b] \times [c, d] \rightarrow R_F$ is said to be continuous in $x_0 \in [a, b], y_0 \in [c, d]$ if for each $\epsilon > 0$ there exist $\delta > 0$ such that $D(f(x, y), f(x_0, y_0)) < \epsilon$, whenever $x \in [a, b], y \in [c, d]$ and $\sqrt{(x - x_0)^2 + (y - y_0)^2} < \delta$. We say that f is fuzzy continuous on $[a, b] \times [c, d]$ if f is continuous at each $(x_0, y_0) \in [a, b] \times [c, d]$.

The space of all such functions is denoted by $C_F([a, b] \times [c, d])$.

Lemma 2.1 [40] If $f, g : [a, b] \times [c, d] \rightarrow R_F$ are fuzzy continuous functions, then the function $F : [a, b] \times [c, d] \rightarrow R^+$ defined by

$$F(x, y) = D(f(x, y), g(x, y))$$

is continuous on $[a, b]$, and $[c, d] \rightarrow R^+$ defined by $F(x, y) = D(f(x, y), g(x, y))$ is continuous on $[a, b]$, and

$$\begin{aligned} D((FR) \int_c^d \int_a^b f(x, y) dx dy, (FR) \int_c^d \int_a^b g(x, y) dx dy) \\ \leq \int_c^d \int_a^b D(f(x, y), g(x, y)) dx dy. \end{aligned}$$

Definition 2.8 . Let $f : [a, b] \times [c, d] \rightarrow R_F$. One call f a uniformly continuous fuzzy real number valued function, if and only if for any $\epsilon > 0$ there exists $\delta > 0$ whenever $\sqrt{(x - s)^2 + (y - t)^2} \leq \delta; x, s \in [a, b], y, t \in [c, d]$, implies that $D(f(x, y)f(s, t)) \leq \epsilon$. one denotes it as $f \in C_F^U([a, b])$.

Definition 2.9 [39] A function $f : [a, b] \times [c, d] \rightarrow R_F$ is said to be bounded if there exist M such that $\|f(x, y)\|_F \leq M$ for any $(x, y) \in [a, b] \times [c, d]$, where $\|f(x, y)\|_F = D(f(x, y), 0)$.

Corollary 2.1 [39] If $f \in C_F([a, b] \times [c, d])$, its definite integral exists [42], furthermore

1. $(FR) \int_c^d \int_a^b f(x, y, r) dx dy = \int_c^d \int_a^b \underline{f}(x, y, r) dx dy,$
2. $(FR) \overline{\int_c^d \int_a^b f(x, y, r) dx dy} = \int_c^d \int_a^b \bar{f}(x, y, r) dx dy$

Remark 2.1 Consider two-dimensional fuzzy Fredholm integral equation of the second kind

$$f(x, y, r) = g(x, y, r) + \lambda \int_c^d \int_a^b K(x, y, s, t) f(s, t, r) ds dt$$

In order to design numerical scheme for solving above equation, we first replace it by the system

$$\underline{f}(x, y, r) = \underline{g}(x, y, r) + \lambda \int_c^d \int_a^b \underline{K}(x, y, s, t) \underline{f}(s, t, r) ds dt$$

$$\bar{f}(x, y, r) = \bar{g}(x, y, r) + \lambda \int_c^d \int_a^b \bar{K}(x, y, s, t) \bar{f}(s, t, r) ds dt$$

where

$$\underline{K}(x, y, s, t) \underline{f}(s, t, r) = \begin{cases} K(x, y, s, t) \underline{f}(s, t, r) & K(x, y, s, t) \geq 0 \\ K(x, y, s, t) \bar{f}(s, t, r) & K(x, y, s, t) < 0 \end{cases}$$

$$\bar{K}(x, y, s, t) \bar{f}(s, t, r) = \begin{cases} K(x, y, s, t) \bar{f}(s, t, r) & K(x, y, s, t) \geq 0 \\ K(x, y, s, t) \underline{f}(s, t, r) & K(x, y, s, t) < 0 \end{cases}$$

Corollary 2.2 [40] If $f, g \in R_F$ are Henstock integrable mappings on $[a, b] \times [c, d]$ and if $D(f(x, y), g(x, y))$ is Lebesgue integrable, then

$$\begin{aligned} D((FH) \int_c^d \int_a^b f(x, y) dx dy, (FH) \int_c^d \int_a^b g(x, y) dx dy) \\ \leq (L) \int_c^d \int_a^b D(f(x, y), g(x, y)) dx dy. \end{aligned}$$

Definition 2.10 [3] Let $f, g : [a, b] \times [c, d] \rightarrow R_F$ be a bounded function, then function

$$\begin{aligned} \omega_{[a, b] \times [c, d]}(f, 0) : R^+ \cup \{0\} \rightarrow R^+, \\ \omega_{[a, b] \times [c, d]} = \sup\{D(f(x, y), f(s, t)) \mid x, s \in [a, b]; y, t \in [c, d], \sqrt{(x - s)^2 + (y - t)^2} \leq \delta\} \end{aligned}$$

where R^+ is the set of positive real numbers, is called the modulus of continuity of f on $[a, b] \times [c, d]$.

Some properties of the modulus of continuity are presented below: [7]

1. $D(f(x, y), f(s, t)) \leq$

$$\omega_{[a,b] \times [c,d]}(f, \sqrt{(x-s)^2 + (y-t)^2})$$

for any $x, s \in [a, b]$ and $y, t \in [c, d]$;

2. $\omega_{[a,b] \times [c,d]}(f, \delta)$ is increasing function of δ ,

3. $\omega_{[a,b] \times [c,d]}(f, 0) = 0$,

4. $\omega_{[a,b] \times [c,d]}(f, \delta_1 + \delta_2) \leq \omega_{[a,b] \times [c,d]}(f, \delta_1) + \omega_{[a,b] \times [c,d]}(f, \delta_2)$ for any $\delta_1, \delta_2 \geq 0$ and $f : [a, b] \times [c, d] \rightarrow R_F$,

5. $\omega_{[a,b] \times [c,d]}(f, n\delta) \leq n\omega_{[a,b] \times [c,d]}(f, \delta)$ for any $\delta \geq 0, n \in N$, and $f : [a, b] \times [c, d] \rightarrow R_F$.

6. $\omega_{[a,b] \times [c,d]}(f, \lambda\delta) \leq ([\lambda] + 1)\omega_{[a,b] \times [c,d]}(f, \delta)$ for any $\delta, \lambda \geq 0$, and any $f : [a, b] \times [c, d] \rightarrow R_F$, where $[\cdot]$ is the ceiling of the number.

In [2], the following theorem is proved.

Corollary 2.3 [2] *Let $f \in C_F([a, b] \times [c, d])$ and the scaling function $\varphi(x, y)$ a real-valued bounded function with it supp*

$\varphi(x, y) \subseteq [-\alpha, \alpha] \times [-\beta, \beta], 0 < \alpha < +\infty, 0 < \beta < +\infty, \varphi(x, y) \geq 0$ such that

$$\sum_{j=-\infty}^{+\infty} \sum_{i=-\infty}^{+\infty} \varphi(x-i)\varphi(y-j) = 1$$

on $[a, b] \times [c, d]$. For $k \in Z^+, x, y \in [a, b] \times [c, d]$, put

$$(B_k f)(x, y) = \sum_{j=-\infty}^{+\infty} \sum_{i=-\infty}^{+\infty} f\left(\frac{i}{2^k}, \frac{j}{2^k}\right) \otimes \varphi(2^k x - i)\varphi(2^k y - j) \tag{2.2}$$

which is a fuzzy-wavelet-like operator. Then

$$D((B_k f)(x, y), f(x, y)) \leq \omega_{[a,b] \times [c,d]}(f, \frac{\alpha}{2^k}, \frac{\beta}{2^k})$$

$$D^*((B_k f), f) \leq \omega_{[a,b] \times [c,d]}(f, \frac{\alpha}{2^k}, \frac{\beta}{2^k})$$

for all $x, y \in R$ and $k \in Z^+$. If $f \in C_F^U([a, b] \times [c, d])$, then as $k \rightarrow +\infty$ one gets $\omega_{[a,b] \times [c,d]}(f, \frac{\alpha}{2^k}, \frac{\beta}{2^k}) \rightarrow 0$ and $\lim_{k \rightarrow +\infty} B_k f = f$, point wise and uniformly with rates.

3 Solving 2FFIE of the second kind

Here, we use fuzzy wavelet like operator defined by (2.2) due to approximate solution of equation (1.1). To do this, we approximate the solution of (1.1) by (2.2). So, by substituting (2.2) in (1.1) we conclude that

$$\sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} f\left(\frac{i}{2^k}, \frac{j}{2^k}\right) \otimes \varphi(2^k x - i)\varphi(2^k y - j) \cong g(x, y) + \lambda \int_c^d \int_a^b K(x, y, s, t) \otimes \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} f\left(\frac{i}{2^k}, \frac{j}{2^k}\right) \otimes (\varphi(2^k s - i)\varphi(2^k t - j)) ds dt \tag{*}$$

By using $2^k s - i = u$ and $2^k t - j = v$, we get

$$(*) \cong g(x, y) +$$

$$\frac{\lambda}{2^{2k}} \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} f\left(\frac{i}{2^k}, \frac{j}{2^k}\right) \otimes$$

$$\int_{2^k c - j}^{2^k d - j} \int_{2^k a - i}^{2^k b - i} K\left(x, y, \frac{u+i}{2^k}, \frac{v+j}{2^k}\right) \varphi(u)\varphi(v) du dv \tag{3.3}$$

Now, by using the following scaling function [38]

$$\varphi(x, y) = \begin{cases} 1 & -\frac{1}{2} \leq x, y \leq \frac{1}{2} \\ 0 & \text{otherwise} \end{cases} \tag{3.4}$$

in (3.3), we conclude that

$$\sum_{j=2^k c}^{2^k d - 1} \sum_{i=2^k a}^{2^k b - 1} f\left(\frac{i}{2^k}, \frac{j}{2^k}\right) \otimes \varphi(2^k x - i)\varphi(2^k y - j) \cong g(x, y) + \frac{\lambda}{2^{2k}} \sum_{j=2^k c}^{2^k d - 1} \sum_{i=2^k a}^{2^k b - 1} f\left(\frac{i}{2^k}, \frac{j}{2^k}\right) \otimes \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} K\left(x, y, \frac{u+i}{2^k}, \frac{v+j}{2^k}\right) \varphi(u)\varphi(v) du dv \tag{3.5}$$

For fixed k , suppose that

$$A_{i,j,k}(x,y) = \frac{\lambda}{2^{2k}} \int_{-\frac{1}{2}}^{\frac{1}{2}} \int_{-\frac{1}{2}}^{\frac{1}{2}} K(x,y, \frac{u+i}{2^k}, \frac{v+j}{2^k}) \varphi(u)\varphi(v)dudv,$$

$$i = 2^k a, 2^k a + 1, \dots, 2^k b - 1,$$

$$j = 2^k c, 2^k c + 1, \dots, 2^k d - 1$$

Clearly, we can write (3.5) in the following form

$$\sum_{j=2^{2k}c}^{2^k d - 1} \sum_{i=2^{2k}a}^{2^k b - 1} f(\frac{i}{2^k}, \frac{j}{2^k}) \otimes \varphi(2^k x_p - i)\varphi(2^k y_q - j)$$

$$\cong g(x_p, y_q) \oplus \sum_{j=2^{2k}c}^{2^k d - 1} \sum_{i=2^{2k}a}^{2^k b - 1} f(\frac{i}{2^k}, \frac{j}{2^k}) \otimes A_{i,j,k}(x_p, y_q)$$

(3.6)

Now, we use collocation points, $x_p = \frac{p}{2^k}, y_q = \frac{q}{2^k}$, $p = 2^k a, 2^k a + 1, \dots, 2^k b - 1, q = 2^k c, 2^k c + 1 \dots 2^k d - 1$ in (3.6). We have:

$$\sum_{j=2^{2k}c}^{2^k d - 1} \sum_{i=2^{2k}a}^{2^k b - 1} f(\frac{i}{2^k}, \frac{j}{2^k}) \otimes \varphi(2^k x_p - i)\varphi(2^k y_q - j)$$

$$= g(x_p, y_q) \oplus \sum_{j=2^{2k}c}^{2^k d - 1} \sum_{i=2^{2k}a}^{2^k b - 1} f(\frac{i}{2^k}, \frac{j}{2^k}) \otimes A_{i,j,k}(x_p, y_q)$$

$$p = 2^k a, 2^k a + 1, \dots, 2^k b - 1$$

$$q = 2^k c, 2^k c + 1, \dots, 2^k d - 1$$

Thus, we obtain the following $2n \times 2n, n = 2^k(b-a)$, fuzzy linear system of equations:

$$C \otimes X = Y \oplus B \otimes X,$$

where

$$X = (f(a, c), f(a + \frac{1}{2^k}, c + \frac{1}{2^k}), \dots, f(b - \frac{1}{2^k}, d - \frac{1}{2^k}))^t$$

$$Y = (g(a, c), g(a + \frac{1}{2^k}, c + \frac{1}{2^k}), \dots, g(b - \frac{1}{2^k}, d - \frac{1}{2^k}))^t,$$

$$B = (B_{i,j})_{2n \times 2n}, B_{ij} = A_{i,j,k}(x_p, y_q),$$

$$C = (c_{i,j})_{2n \times 2n}, c_{ij} = \varphi(2^k x_p - i)\varphi(2^k y_q - j)$$

Here, we suppose that $b - a \in N, d - c \in N$, where N is set of natural numbers. Clearly, the above system is a dual fuzzy linear system. For solving this system, one can refer to [37]. Finally, by solving this system and using Eq (2.2), we can present the approximate solution of (1.1).

4 Error Estimation

Corollary 4.1 Consider linear 2FFIE of the second kind as follows:

$$f(x,y) = (g(x,y) \oplus \lambda \int_c^d \int_a^b K(x,y,s,t) \otimes f(s,t)dsdt,$$

where $K(x,y,s,t)$ is an arbitrary continues kernel function having same sign in the square $a \leq x, s \leq b, c \leq y, t \leq d$, and $g(x,y) \neq 0$ is a continuous fuzzy function over $a \leq x \leq b, c \leq y \leq d$. Under hypothesis of Theorem(2.3), we have:

$$D(f(x,y), (B_k f)(x,y)) \leq M|\lambda|(b-a)(d-c)$$

$$\omega_{[a,b] \times [c,d]}(f, \frac{\alpha}{2^k}, \frac{\beta}{2^k})$$

where $M = \max|K(x,y,s,t)|, a \leq x, s \leq b$ and $c \leq y, t \leq d$, and

$$(B_k f)(x,y) = \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} f(\frac{i}{2^k}, \frac{j}{2^k}) \otimes$$

$$(\varphi(2^k x - i)\varphi(2^k y - j))$$

is the approximate solution of equation (1.1). **Proof.** We would like to estimate

$$D(f(x,y), (B_k f)(x,y)) = D(g(x,y) \oplus$$

$$\lambda \int_c^d \int_a^b K(x,y,s,t) \otimes f(s,t)dsdt,$$

$$g(x,y) \oplus \lambda \int_c^d \int_a^b K(x,y,s,t) \otimes (B_k f)(s,t)dsdt) =$$

$$D(\lambda \int_c^d \int_a^b K(x,y,s,t) \otimes f(s,t)dsdt,$$

$$\lambda \int_c^d \int_a^b K(x,y,s,t) \otimes (B_k f)(s,t)dsdt)$$

$$\leq |\lambda| \int_c^d \int_a^b |K(x,y,s,t)| \cdot D(f(s,t), (B_k f)(s,t))dsdt$$

$$\leq |\lambda| \int_c^d \int_a^b M \cdot D^*(f, B_k f)dsdt \leq$$

$$M|\lambda|(b-a)(d-c)\omega_{[a,b] \times [c,d]}(f, \frac{\alpha}{2^k}, \frac{\beta}{2^k}) \blacksquare$$

It is obvious that

$$(\overline{B_k f})(x,y) = \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \overline{f}(\frac{i}{2^k}, \frac{j}{2^k}) \otimes$$

$$\varphi(2^k x - i)\varphi(2^k y - j)$$

$$(\underline{B_k f})(x,y) = \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \underline{f}(\frac{i}{2^k}, \frac{j}{2^k}) \otimes$$

$$\varphi(2^k x - i)\varphi(2^k y - j).$$

Now, we consider two cases as follows:

- (a) $K(x, y, s, t) \geq 0$,
- (b) $K(x, y, s, t) < 0$.

In case (a), we have:

$$\begin{aligned} \bar{e} &= \bar{f}(x, y) - \overline{B_k f}(x, y) = [\bar{g}(x, y) + \\ &\lambda \int_c^d \int_a^b K(x, y, s, t) \otimes \bar{f}(s, t) ds dt] - \\ &[\bar{g}(x, y) + \lambda \int_c^d \int_a^b K(x, y, s, t) (\overline{B_k f})(s, t) ds dt] \\ &= \lambda \int_c^d \int_a^b K(x, y, s, t) \otimes \bar{f}(s, t) ds dt - \\ &\lambda \int_c^d \int_a^b K(x, y, s, t) (\overline{B_k f})(s, t) ds dt \\ &= \lambda \int_c^d \int_a^b K(x, y, s, t) [\bar{f}(s, t) - (\overline{B_k f})(s, t)] ds dt \end{aligned}$$

similary we have:

$$\begin{aligned} \underline{e} &= \underline{f}(s, t) - (\underline{B_k f})(s, t) \\ \underline{g}(x, y) &+ \lambda \int_c^d \int_a^b K(x, y, s, t) \underline{f}(s, t) ds dt \\ -[\underline{g}(x, y) &+ \lambda \int_c^d \int_a^b K(x, y, s, t) (\underline{B_k f})(s, t) ds dt]. \\ &= \lambda \int_c^d \int_a^b K(x, y, s, t) \underline{f}(s, t) ds dt \\ &- \lambda \int_c^d \int_a^b K(x, y, s, t) (\underline{B_k f})(s, t) ds dt \\ &= \lambda \int_c^d \int_a^b K(x, y, s, t) [\underline{f}(s, t) - (\underline{B_k f})(s, t)] ds dt \end{aligned}$$

Hence

$$\begin{aligned} e &= \bar{e} - \underline{e} = \lambda \int_c^d \int_a^b K(x, y, s, t) [\bar{f}(s, t) - (\overline{B_k f})(s, t)] ds dt \\ &\lambda \int_c^d \int_a^b K(x, y, s, t) [\underline{f}(s, t) - (\underline{B_k f})(s, t)] ds dt \\ &= \lambda \int_c^d \int_a^b K(x, y, s, t) ([\bar{f}(s, t) - (\overline{B_k f})(s, t)] \\ &+ [\underline{f}(s, t) - (\underline{B_k f})(s, t)]) ds dt, \end{aligned}$$

and

$$\begin{aligned} \|e\| &= \|\bar{e} + \underline{e}\| \leq |\lambda| \int_c^d \int_a^b \|K(x, y, s, t)\| \|[\bar{f}(s, t) - (\overline{B_k f})(s, t)]\| \\ &+ \|\underline{f}(s, t) - (\underline{B_k f})(s, t)\| ds dt. \end{aligned}$$

Since $K(x, y, s, t)$ is an arbitrary continues kernel function over the square $a \leq x, s \leq b, c \leq y, t \leq d$, there exist $M > 0$ such that $M = \max|K(x, y, s, t)|$. So, we can write

$$\begin{aligned} \|e\| &\leq |\lambda| M \int_c^d \int_a^b \|[\bar{f}(s, t) - (\overline{B_k f})(s, t)]\| \\ &+ \|\underline{f}(s, t) - (\underline{B_k f})(s, t)\| ds dt. \end{aligned} \tag{4.7}$$

On the other hand, we have:

$$\begin{aligned} \bar{f}(s, t) - \overline{B_k f}(s, t) &= \bar{f}(s, t) \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \varphi(2^k x - i) \varphi(2^k y - j) - \\ &\sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \bar{f}(\frac{i}{2^k}, \frac{j}{2^k}) \otimes \varphi(2^k x - i) \varphi(2^k y - j) \\ &= \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} [\bar{f}(s, t) - \bar{f}(\frac{i}{2^k}, \frac{j}{2^k})] \varphi(2^k x - i) \varphi(2^k y - j), \end{aligned}$$

Therefore

$$\begin{aligned} \|\bar{f}(s, t) - (\overline{B_k f})(s, t)\| &= \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} [\bar{f}(s, t) - \bar{f}(\frac{i}{2^k}, \frac{j}{2^k})] \\ &\varphi(2^k x - i) \varphi(2^k y - j) \| \\ &\leq \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \|\bar{f}(s, t) - \bar{f}(\frac{i}{2^k}, \frac{j}{2^k})\| \varphi(2^k x - i) \varphi(2^k y - j) \\ &\leq \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \omega[\bar{f} \|x - \frac{i}{2^k}, y - \frac{j}{2^k}\|] \\ &\varphi(2^k x - i) \varphi(2^k y - j), \quad (**) \end{aligned}$$

Notice that, under hypotheses of Theorem (2.3), we conclude that

$$\begin{aligned} -\alpha \leq 2^k x - i \leq \alpha &\Rightarrow -\frac{\alpha}{2^k} \leq x - \frac{i}{2^k} \leq \frac{\alpha}{2^k} \\ &\Rightarrow |x - \frac{i}{2^k}| \leq \frac{\alpha}{2^k}, \\ -\beta \leq 2^k x - i \leq \beta &\Rightarrow -\frac{\beta}{2^k} \leq x - \frac{i}{2^k} \leq \frac{\beta}{2^k} \\ &\Rightarrow |x - \frac{i}{2^k}| \leq \frac{\beta}{2^k}. \end{aligned}$$

Hence

$$(**) \leq \sum_{j=-\infty}^{\infty} \sum_{i=-\infty}^{\infty} \varphi(2^k x - i) \varphi(2^k y - j)$$

$$\omega_{[a,b] \times [c,d]}(\bar{f}, \frac{\alpha}{2^k}, \frac{\beta}{2^k}) = \omega_{[a,b] \times [c,d]}(\bar{f}, \frac{\alpha}{2^k}, \frac{\beta}{2^k})$$

Therefore, as $k \rightarrow +\infty$ we get

$$\|\bar{f}(s, t) - (\overline{B_k f})(s, t)\| \rightarrow 0$$

Similarly, we conclude

$$\|\underline{f}(s, t) - (\underline{B_k f})(s, t)\| \rightarrow 0, \text{ as } k \rightarrow +\infty.$$

So we obtain that

$$\|e\| = \|\bar{e} + \underline{e}\| \rightarrow 0,$$

as $k \rightarrow +\infty$.

In case (b) we have $K(x, y, s, t) < 0$. So

$$\begin{aligned} \bar{e} &= [\bar{g}(x, y) + \lambda \int_c^d \int_a^b K(x, y, s, t) \underline{f}(s, t) ds dt \\ &- [\bar{g}(x, y) + \lambda \int_c^d \int_a^b K(x, y, s, t) (\underline{B}_k f)(s, t) ds dt] \\ &= +\lambda \int_c^d \int_a^b K(x, y, s, t) (\underline{f}(s, t) - \underline{B}_k f)(s, t) ds dt \end{aligned}$$

and similarly way, we have:

$$\underline{e} = \lambda \int_c^d \int_a^b K(x, y, s, t) [\bar{f}(s, t) - (\overline{B}_k f)(s, t)] ds dt$$

Hence

$$\|e\| = \|\bar{e} + \underline{e}\|$$

$$\begin{aligned} &\leq |\lambda| M \int_c^d \int_a^b \|K(x, y, s, t)\| \|[\underline{f}(s, t) - (\underline{B}_k f)(s, t)]\| + \\ &\quad \|[\bar{f}(s, t) - (\overline{B}_k f)(s, t)]\| ds dt \\ &\leq |\lambda| M (b-a)(d-c) [\omega_{[a,b] \times [c,d]}(\underline{f}, \frac{\alpha}{2^k}, \frac{\beta}{b^k}) \\ &\quad + \omega_{[a,b] \times [c,d]}(\bar{f}, \frac{\alpha}{2^k}, \frac{\beta}{2^k})] \end{aligned}$$

Clearly, we have:

$$\|e\| = \|\bar{e} + \underline{e}\| \rightarrow 0, \text{ as } k \rightarrow +\infty.$$

5 Numerical examples

To illustrate the efficiency of the presented method in Section , we give two examples. Also, we compare the numerical solutions obtained by using the proposed method with the exact solutions. Trough this section, we suppose that

$$\varphi(x, y) = \begin{cases} 1 & -\frac{1}{2} \leq x, y \leq \frac{1}{2} \\ 0 & \text{otherwise} \end{cases}$$

Example 5.1 Consider the following linear 2FFIE of the second kind

$$\underline{g}(x, y, r) = (x \sin \frac{y}{2})(r^2 + r)$$

$$\bar{g}(x, y) = (x \sin \frac{y}{2})(4 - r^3 - r)$$

$$K(x, y, s, t) = x^2 y s; 0 \leq x, y, s, t \leq 1$$

Also, let $a = 0, b = 1$. The exact solution of this example is given by

$$\underline{f}(x, y, r) = [(x \sin \frac{y}{2}) - \frac{16}{21} (\cos \frac{1}{2} - 1) . x^2 y](r^2 + r)$$

$$\bar{f}(x, y, r) = [(x \sin \frac{y}{2}) - \frac{16}{21} (\cos \frac{1}{2} - 1) . x^2 y](4 - r^3 - r)$$

By using the proposed method in Section 5, we can present the approximate solution for this example. To compare the numerical results with the exact solution for different values of x, y and k , see Tables 1-3.

Example 5.2 Consider the following linear 2FFIE of the second kind

$$\underline{g}(x, y, r) = r(\frac{1}{3}r + \frac{8}{3})(1 + x + y - \frac{7}{12}xy)$$

$$\bar{g}(x, y) = (2r^2 - 4r + 5)(1 + x + y - \frac{7}{12}xy)$$

$$K(x, y, s, t) = xyst; \quad 0 \leq x, y, s, t \leq 1$$

Also, let $a = 0, b = 1$. The exact solution of this example is given by

$$\underline{f}(x, y, r) = r(\frac{1}{3}r + \frac{8}{3})(x + y + 1)$$

$$\bar{f}(x, y, r) = (2r^2 - 4r + 5)(x + y + 1)$$

By using the proposed method in Section 5, we can present the approximate solution for this example. To compare the numerical results with the exact solution for different values of x, y and k , see Tables 4-5.

6 Conclusion

To approximate the solution of 2FFIE of the second kind, a new approach based on fuzzy wavelet like operator via a real-valued scaling function and collocation method is proposed. Convergence analysis of the proposed method is investigated by using the modulus of continuity in one theorem. To illustrate the efficiency of the presented method, two examples are given. Comparing the numerical solutions with the exact solution show that the proposed method can be a suitable method for solving 2FFIE of the second numerically .

Table 1: Numerical results for Example 5.1 for $k = 2, x = \frac{1}{4}, y = \frac{1}{2}$

r	$ f(x, y) - (B_k f)(x, y) $	$ \bar{f}(x, y) - \overline{B_k f}(x, y) $
0.1	0.000869398	0.0308162
0.2	0.00189687	0.0299705
0.3	0.00189687	0.02903
0.4	0.00442603	0.0279472
0.5	0.00592771	0.0266747
0.6	0.00758747	0.0251651
0.7	0.00940531	0.023371
0.8	0.0113812	0.0212449
0.9	0.0135152	0.0187395
1.0	0.0158072	0.0158072

Table 2: Numerical result for Example 5.1 with $k = 3, x = \frac{1}{8}, y = \frac{2}{8}$

r	$ f(x, y) - (B_k f)(x, y) $	$ \bar{f}(x, y) - \overline{B_k f}(x, y) $
0.1	0.0000430158	0.00152471
0.2	0.0000938526	0.00148287
0.3	0.000152511	0.00143634
0.4	0.00021899	0.00138276
0.5	0.00029329	0.0013198
0.6	0.000375411	0.00124511
0.7	0.000465353	0.00115634
0.8	0.000563116	0.00103115
0.9	0.0006687	0.000927186
1.0	0.000782105	0.000782105

Table 3: Numerical results for Example 5.1 for $k = 4, x = \frac{1}{16}, y = \frac{1}{8}$

r	$ f(x, y) - (B_k f)(x, y) $	$ \bar{f}(x, y) - \overline{B_k f}(x, y) $
0.1	5.0409210^{-6}	0.00152471
0.2	0.0000109984	0.00148287
0.3	0.0000178724	0.00143634
0.4	0.0000256629	0.00138276
0.5	0.0000343699	0.0013198
0.6	0.0000439935	0.00124511
0.7	0.0000545336	0.00115634
0.8	0.0000783634	0.00103115
0.9	0.0006687	0.000927186
1.0	0.0000916531	0.000782105

References

- [1] S. Abbasbandy, E. Babolian, M. Alavi, *Numerical method for solving linear fredholm fuzzy integral equations of the second kind*, Chaos Solutions an Fractals 31 (2007) 138-146.
- [2] R. Ezzati, F. Mokhtarnejad, N. Hassasi, *Some fuzzy-wavelet-like operators and their convergence*, Mathematical Problems in Engineering (2013), Article ID 832831, 10 pages, <http://dx.doi.org/10.1155/2013/832831/>.
- [3] G. A. Anastassiou, *Fuzzy mathematics: Approximation theory*, Springer-Verlag Berlin Heidelberg (2010).
- [4] E. Babolian, H. Sadeghi Goghary, S. Ab-

Table 4: Numerical results for Example 5.2 for $k = 2, x = \frac{1}{4}, y = \frac{1}{4}$

r	$ f(x, y) - (B_k f)(x, y) $	$ \bar{f}(x, y) - \overline{B_k f}(x, y) $
0.1	0.0235411	0.402814
0.2	0.0476634	0.37317
0.3	0.072367	0.347013
0.4	0.0976519	0.324344
0.5	0.123518	0.305162
0.6	0.149965	0.289468
0.7	0.176994	0.277262
0.8	0.204604	0.268543
0.9	0.232795	0.263311
1.0	0.261568	0.261568

Table 5: Numerical results for Example 5.2 for $k = 3, x = \frac{1}{8}, y = \frac{1}{8}$

r	$ f(x, y) - (B_k f)(x, y) $	$ \bar{f}(x, y) - \overline{B_k f}(x, y) $
0.1	0.00283428	0.0484977
0.2	0.00573855	0.0449286
0.3	0.00871279	0.0417794
0.4	0.117557	0.0390501
0.5	0.0148712	0.0367407
0.6	0.0180554	0.0348512
0.7	0.0213096	0.0333815
0.8	0.0246338	
0.9	0.0280279	0.031702
1.0	0.031492	0.031492

basbandy, *Numerical solution of linear Fredholm fuzzy integral equations of the second kind by Adomian method*, Appl. Math. Comput. 161 (2006) 733-744 .

[5] K. Balachandran, K. Kanagarajan, *Existence of solutions general nonlinear fuzzy Volterra-Fredholm integral equations*, Appl. Math. Stochastic Anal. 3 (2005) 333-343.

[6] K. Balachandran, P. Prakash, *Existence of solutions of nonlinear fuzzy Volterra-Fredholm integral equations*, Indian J. Pure Appl. Math. 33 (2002) 329-343.

[7] B. Bede, S. G. Gal, *Quadrature rules for integrals of fuzzy-number-valued functions*, Fuzzy Sets and Systems 145 (2004) 359-380.

[8] A. M. Bica, *Error estimation in the approximation of the solution of nonlinear fuzzy Fredholm integral equations*, Information Sciences 178 (2008) 1279-1292.

[9] A. M. Bica, C. Popescu, *Numerical solutions of the nonlinear fuzzy Hammerstein-Volterra delay integral equations*, Information Sciences 223 (2013) 236-255.

[10] A. M. Bica, *One-sided fuzzy numbers and applications to integral equations form epidemiology*, Fuzzy Sets and Systems 219 (2013) 27-48.

[11] D. Dubois, H. Prade, *Towards fuzzy differential calculus*, Fuzzy Sets and Systems 8 (1982) 1-7.

[12] R. Ezzati, S. Ziari, *Numerical solution and error estimation of fuzzy Fredholm integral equation using fuzzy Bernstein polynomials*, Aust. J. Basic Appl. Sci. 5 (2011) 2072-2082.

[13] M. A . Fariborzi Araghi, N. Parandin, *Numerical solution of fuzzy Fredholm integral equations by the Lagrange interpolation based on the extension principle*, Soft Computing 15 (2011) 2449-2456.

- [14] M. Friedman, M. Ma, A. Kandel, Numerical methods for calculating the fuzzy integral, *Fuzzy Sets and Systems* 83 (1996) 57-62.
- [15] M. Friedman, M. Ma, A. Kandel, *On fuzzy integral equations*, *Fund. Inform.* 37 (1999) 89-99.
- [16] M. Friedman, M. Ma, A. Kandel, *Solutions to fuzzy integral equations with arbitrary kernels*, *International Journal of Approximate Reasoning* 20 (1999) 249-262.
- [17] M. Friedman, M. Ma, A. Kandel, *Numerical solutions of fuzzy differential and integral equations*, *Fuzzy Sets and Systems* 106 (1999) 35-48.
- [18] S. Ziari, R. Ezzati, S. Abbasbandy, *Numerical solution of linear fuzzy Fredholm integral equations of the second kind using fuzzy Haar wavelets*, *Commun. Comput. Inf. Sci.* 299 (3) (2012) 79-89.
- [19] D. N. Georgiou, I. E. Kougias, *Bounded solutions for fuzzy integral equations*, *Int. J. Math. Math. Sci.* 31 (2002) 109-114.
- [20] R. Goetschel, W. Voxman, *Elementary fuzzy calculus*, *Fuzzy Sets and Systems* 18 (1986) 31-43. O. Kaleva, *Fuzzy differential equations*, *Fuzzy Sets and Systems* 24 (1987) 301-317.
- [21] J. Mordeson, W. Newman, *Fuzzy integral equations*, *Inform. Sci.* 87 (1995) 215-229.
- [22] S. Nanda, *On integration of fuzzy mappings*, *Fuzzy Sets and Systems* 32 (1989) 95-101.
- [23] J. J. Nieto, R. Rodriguez-Lopez, *Bounded solutions for fuzzy differential and integral equations*, *Chaos Solitons Fractals* 27 (2006) 1376-1386.
- [24] N. Parandin, M. A. Fariborzi Araghi, *The numerical solution of linear fuzzy Fredholm integral equations of the second kind by using finite and divided differences methods*, *Soft Computing* 15 (2010) 729-741.
- [25] J. Y. Park, S. Y. Lee, J. U. Jeong, *The approximate solution of fuzzy functional integral equations*, *Fuzzy Sets and Systems* 110 (2000) 79-90.
- [26] J. Y. Park, J. U. Jeong, *On the existence and uniqueness of solutions of fuzzy Volterra-Fredholm integral equations*, *Fuzzy Sets and Systems* 115 (2000) 425-431.
- [27] J. Y. Park, J. U. Jeong, *A note on fuzzy integral equations*, *Fuzzy Sets Systems* 108 (1999) 193-200.
- [28] J. Y. Park, H. K. Han, *Existence and uniqueness theorem for a solution of fuzzy Volterra integral equations*, *Fuzzy Sets Systems* 105 (1999) 481-488.
- [29] A. Jafarian, S. Measoomy Nia, S. Tavan, *A numerical scheme to solve fuzzy linear Volterra integral equations system*, *Journal of Applied Mathematics* (2012) Article ID 216923, 17 pages <http://dx.doi.org/doi:10.1155/2012/216923/>.
- [30] H. Sadeghi Goghary, M. Sadeghi Goghary, *Two computational methods for solving linear Fredholm fuzzy integral equations of the second kind by Adomian method*, *Appl. Math. Comput.* 161 (2005) 733-744.
- [31] M. Shafiee, S. Abbasbandy, T. Allahviranloo, *Predictor-corrector method for nonlinear fuzzy Volterra integral equations*, *Aust. J. Basic Appl. Sci.* 5 (2011) 2865-2874.
- [32] C. Wu, Z. Gong, *On Henstock integral of fuzzy-number-valued functions*, *Fuzzy Sets and Systems* 120 (2001) 523-532.
- [33] H. C. Wu, *The fuzzy Riemann integral and its numerical integration*, *Fuzzy Sets and Systems* 110 (2000) 1-25.
- [34] M. Friedman, M. Ming, A. Kandel, *Fuzzy linear systems*, *Fuzzy Sets and Systems* 96 (1998) 201-209.
- [35] G. A. Anastassiou, *Fuzzy wavelet type operators*, *Nonlinear Functional Analysis and Applications* 9 (2004) 251-269.
- [36] A. Rivaz, F. Yousefi, *Modified Homotopy Perturbation Method for Solving Two-Dimensional Fuzzy Fredholm Integral Equation*, *Fuzzy Sets and Systems* 25 (2012) 591-602.

- [37] S. M. Sadatrasoul, R. Ezzati, *Quadrature Rules and Iterative Method for Numerical Solution of Two-Dimensional Fuzzy Integral Equations*, (2014) Article ID 413570.
- [38] F. Mokhtarnejad, R. Ezzati, *Existence and uniqueness of the solution of fuzzy-valued integral equations of mixed type*, Iranian Journal of Fuzzy Systems 12 (2015) 87-94.
- [39] R. Ezzati, S. Ziari, *Numerical Solution of Two-Dimensional Fuzzy Fredholm Integral Equations of the Second kind Using Fuzzy Bivariate Bernestein Polynomials*, Fuzzy Sets and Systems 15 (2013) 84-89.
- [40] S. M. Sadatrasoul, R. Ezzati, *Iterative method for numerical solution of two-dimensional nonlinear fuzzy integral equations*, Fuzzy Sets Syst. 12 (2014) <http://dx.doi.org/doi:10.1016/j.fss.2014.12.008>.
- [41] M. Baghmisheh, R. Ezzati, *Numerical solution of fuzzy Fredholm integral equations of the second kind using hybrid of block-pulse functions and Taylor series*, Advances in Difference Equations (2015) <http://dx.doi.org/doi:10.1186/s13662-015-0389-7/>.
- [42] R. Ezzati, S. Ziari, *Numerical solution of nonlinear fuzzy Fredholm integral equations using iterative method*, Appl. Math. Comput. 225 (2013) 33-42.
- [43] F. Mokhtarnejad, R. Ezzati, *The numerical solution of nonlinear Hammerstein fuzzy integral equations by using fuzzy wavelet like operator*, Journal of Intelligent and Fuzzy Systems 28 (2015) 1617-1628.
- [44] S. M. Sadatrasoul, R. Ezzati, *Numerical solution of two-dimensional nonlinear Hammerstein fuzzy integral equations based on optimal fuzzy quadrature formula*, Journal of Computational and Applied Mathematics 292 (2016) 430-446.



Nader Hassasi was born in Marand, Iran in 1974. He received his B.S. degree in Teaching Mathematics in 1997 from Marand Branch Islamic Azad University, and M.S. in 1999 from Lahijan Branch Islamic Azad University.

He is currently the member of board at the IAU-Malayer Branch where he is a Scientific Club member since (2007). Since 2011 he is Ph.D research student in Numerical Analysis at Islamic Azad University, Karaj Branch, Iran. His main research interest is fuzzy mathematics especially, on numerical solution of fuzzy integral equations.



Reza Ezzati was born in Tabriz, Iran in 1974. He received his B. S. degree in pure mathematics in 1997 and M.S. degree in applied mathematics from Azerbaijan Teacher Education University and Tarbyat Modarres University of Tehran, respectively, and his Ph.D degree in applied mathematics from IAU- Science and Research Branch, Tehran, Iran in 2006. He is an Associate Professor in the Department of Mathematics at Islamic Azad University, Karaj Branch, (Iran). He has published about 95 papers in international journals, and he also is the Associate Editor of Mathematical Sciences (a Springer Open Journal). He is the Member of Editorial Board of The Scientific Journal, Journal of Applied Mathematics. He is a Reviewer in the Journals of Iranian Journal of Fuzzy systems, Information Science (Elsevier), Applied Mathematics Letters (Elsevier), Intelligent and Fuzzy Systems, Applied Mathematical Modelling, Applied Mathematics and Computations, and others. His current interests include fuzzy mathematics, especially, on numerical solution of fuzzy systems, fuzzy integral equations and fuzzy interpolation.

He is an Associate Professor in the Department of Mathematics at Islamic Azad University, Karaj Branch, (Iran). He has published about 95 papers in international journals, and he also is the Associate Editor of Mathematical Sciences (a Springer Open Journal). He is the Member of Editorial Board of The Scientific Journal, Journal of Applied Mathematics. He is a Reviewer in the Journals of Iranian Journal of Fuzzy systems, Information Science (Elsevier), Applied Mathematics Letters (Elsevier), Intelligent and Fuzzy Systems, Applied Mathematical Modelling, Applied Mathematics and Computations, and others. His current interests include fuzzy mathematics, especially, on numerical solution of fuzzy systems, fuzzy integral equations and fuzzy interpolation.