



The spectral iterative method for Solving Fractional-Order Logistic Equation

A. Shoja ^{*†}, E. Babolian [‡], A. R. Vahidi [§]

Received Date: 2014-09-19 Revised Date: 2015-01-16 Accepted Date: 2015-06-19

Abstract

In this paper, a new spectral-iterative method is employed to give approximate solutions of fractional logistic differential equation. This approach is based on combination of two different methods, i.e. the iterative method [35] and the spectral method. The method reduces the differential equation to systems of linear algebraic equations and then the resulting systems are solved by a numerical method. The solutions obtained are compared with Adomian decomposition method and iterative method used in [35] and Adams method [36].

Keywords : Adomian decomposition method (ADM); Iterative method (IM); Spectral method; Fractional logistic equation; Collocation method.

1 Introduction

To describe population growth in a limited environment, Verhulst [28] first presented the classical logistic equation and it has been very popular in population dynamics so far. We can apply the fractional derivative operator on the logistic equation to obtain the fractional order logistic model. Pierre Verhulst published this model in 1838 for the first time [14]. We can describe the continuous logistic model by first order ordinary differential equation. The discrete logistic model is a simple iterative equation which shows the chaotic property in certain regions [11, 29]. There are many variations of the population modeling. To describe the periodic

doubling and chaotic characteristic in dynamical system we can use Verhulst model which is a classic example [11]. This model indicates that the population growth may be restricted by some factors like population density [12, 23].

Many studies are focussed on ordinary and partial fractional equations thanks to their recurrent appearance in different applications in fluid mechanics, viscoelasticity, biology, physics and engineering [13]. Most recently, a large amount of literatures are developed regarding the usage of fractional differential equations in non-linear dynamics. Consequently, the solutions of fractional differential equations of physical interest have been of great importance. We can not find exact solutions for most fractional differential equations, so approximate and numerical techniques are applied [15, 16, 19, 20, 21, 22]. Recently to solve the fractional differential equations several numerical and approximate methods, such as variational iteration method [17], iterative method [35], homotopy perturbation method [24],

*Corresponding author. shoja@riau.ac.ir

[†]Department of Mathematics, Science and Research Branches, Islamic Azad University, Tehran, Iran.

[‡]Department of Mathematics, Science and Research Branches, Islamic Azad University, Tehran, Iran.

[§]Department of Mathematics, Science and Research Branches, Islamic Azad University, Tehran, Iran.

Adomian decomposition method [8], homotopy analysis method and collocation method [18, 26] have been employed.

We consider fractional logistic equation of the following form:

$$\begin{cases} D^\alpha y(x) = \mu y(x)(1 - y(x)), \\ y(0) = y_0. \end{cases} \quad (1.1)$$

where $\mu > 0, x > 0, 0 < \alpha \leq 1$.

The important application of the logistic equation is that it is a model of population growth. The population size at time x is denoted with $y(x)$ and the constant $\mu > 0$ defines the growth rate. Another application of Logistic equation is in medicine, where the logistic differential equation is used to model the growth of tumors. This application can be considered as an extension of the above mentioned use in the frame work of ecology. The existence and the uniqueness of the solution to the proposed problem (1.1) are introduced in details in [6].

In this paper, we describe preliminaries in Sec. 2, in Sec. 3.1 we describe the iterative method and in Sec. 3.2 we give a description of shifted fractional-order Legendre functions. In Sec. 3.3 we use collocation method to obtain the approximate solution for differential equation with initial conditions as a linear combination of Legendre functions. In Sec. 3.4, we describe the new spectral-iterative method (NSIM) which is a combination of two different methods, one iterative and the other spectral. We study the numerical results in Sec. 4 and review the estimation of the errors in Sec. 5.

2 Preliminaries

Definition 2.1 *In order to proceed, we need the following definitions of fractional derivatives and integrals. First, we introduce the Riemann-Liouville definition of fractional integral operator J_a^α .*

Let $\alpha \in R^+$. The operator J_a^α , defined on the usual Lebesgue space $L_1[a, b]$ by

$$J_a^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, \quad (2.2)$$

$$J_a^0 f(x) = f(x),$$

for $a \leq x \leq b$, is called the Riemann-Liouville fractional integral operator of order α .

Properties of the operator J_a^α can be found in [1]. For $f \in L_1[a, b], \alpha, \beta \geq 0$ and $\gamma > -1$, we mention only the following:

$$(1) \quad J_a^\alpha f(x) \text{ exists for almost every } x \in [a, b],$$

$$(2) \quad J_a^\alpha J_a^\beta f(x) = J_a^{\alpha+\beta} f(x),$$

$$(3) \quad J_a^\alpha J_a^\beta f(x) = J_a^\beta J_a^\alpha f(x),$$

$$(4) \quad J_a^\alpha (x-a)^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(\alpha+\gamma+1)} (x-a)^{\alpha+\gamma}.$$

Definition 2.2 *The fractional derivative of $f(x)$ in the Riemann-Liouville sense is defined as*

$$D_a^\alpha f(x) = D^m J_a^{m-\alpha} f(x) = \frac{d^m}{dx^m} \frac{1}{\Gamma(m-\alpha)} \int_a^x (x-t)^{m-\alpha-1} f(t) dt, \quad (2.3)$$

where $m \in N$ and satisfies the relations $m-1 < \alpha \leq m$, and $f \in L_1[a, b]$.

Properties of the operator D_a^α can be found in [1, 4]. For $m-1 < \alpha \leq m, x > a$ and $\gamma > -1$ we mention only the following:

$$(1) \quad D_a^\alpha (x-a)^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(\gamma-\alpha+1)} (x-a)^{\gamma-\alpha},$$

$$(2) \quad D_a^\alpha J_a^\alpha f(x) = f(x).$$

3 New spectral iterative method

	$\alpha = \frac{1}{2}, \mu = \frac{1}{5}, \beta = \frac{1}{4}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$1.6E-07$	0.204
10	$1.0E-13$	0.219
15	$1.8E-20$	0.484
20	$1.6E-27$	0.782

Table 1.

	$\alpha = \frac{1}{4}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$1.2E - 08$	0.250
10	$9.0E - 15$	0.232
15	$3.0E - 21$	0.437
20	$8.0E - 28$	1.484

Table 2.

	$\alpha = \frac{4}{5}, \mu = \frac{1}{2}, \beta = \frac{1}{5}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$3.0E - 06$	0.172
10	$2.0E - 12$	0.266
15	$2.0E - 19$	0.453
20	$6.0E - 27$	0.844

Table 4.

	$\alpha = \frac{3}{10}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$8.0E - 09$	0.187
10	$3.0E - 15$	0.250
15	$6.0E - 22$	0.656
20	$7.0E - 29$	1.531

Table 3.

	$\alpha = \frac{1}{2}, \mu = \frac{1}{5}, \beta = \frac{1}{4}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$5.0E - 08$	1.297
10		failed
15		failed
20		failed

Table 5.

3.1 Iterative method

Consider the following nonlinear differential equation:

$$L[y] + N[y] = f(x), \tag{3.4}$$

where L is a linear operator and N is a nonlinear operator from a Banach space E into E , f is a given function in E and we are looking for $y \in E$ satisfying (3.4).

Daftardar and Jafari [35], suggest that the solution of $y(x)$ be expanded by the infinite series solution

$$y(x) = \sum_{k=0}^{\infty} y_k(x), \tag{3.5}$$

and the nonlinear operator N in Eq. (3.4) is decomposed as follows:

$$N(y) = \sum_{i=0}^{\infty} A_i(y_0, y_1, \dots, y_i), \tag{3.6}$$

where $A_0 = 0$ and A_i are obtained by

$$A_i = N\left(\sum_{k=0}^i y_k\right) - N\left(\sum_{k=0}^{i-1} y_k\right).$$

Substituting (3.5) and (3.6) into (3.4) gives the following recursive scheme:

$$\begin{cases} L[y_0] = f(x), \\ L[y_{i+1}] = -A_i, \quad i = 0, 1, \dots \end{cases} \tag{3.7}$$

We define the $M + 1$ -th term approximation solution as

$$\phi_M(x) = \sum_{i=0}^M y_i(x), \tag{3.8}$$

where, if convergence happen,

$$y(x) = \lim_{M \rightarrow \infty} \phi_M(x).$$

3.2 Shifted fractional-order Legendre function

The Legendre polynomials, denoted by $l_n(x)$, are orthogonal with respect to the weight function $w(x) = 1$ over $I = [-1, 1]$, namely [9],

$$\int_{-1}^1 l_n(x)l_m(x)dx = \frac{2}{2n+1}\delta_{nm},$$

where

$$\delta_{nm} = \begin{cases} 1, & n = m, \\ 0, & O.W. \end{cases}$$

In order to use these polynomials on the interval $[0, 1]$, we define the so-called shifted Legendre polynomials by introducing the change of variable $x = 2t - 1$. Let the shifted Legendre polynomials $l_n(x)$ be denoted by $L_n(t)$. The shifted Legendre polynomials are orthogonal with respect to the weight function $w(t) = 1$ in the interval $[0, 1]$ with the orthogonality property

$$\int_0^1 L_n(t)L_m(t)dt = \frac{2}{2n+1}\delta_{nm}.$$

Then $L_i(t)$ can be obtained as follows:

$$L_{n+1}(t) = \frac{(2n+1)(2t-1)}{n+1}L_n(t) - \frac{n}{n+1}L_{n-1}(t), n = 1, 2, \dots$$

	$\alpha = \frac{1}{4}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$4.0E - 06$	3.891
10		failed
15		failed
20		failed

Table 6.

	$\alpha = \frac{4}{5}, \mu = \frac{1}{2}, \beta = \frac{1}{5}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$2.0E - 06$	4.297
10		failed
15		failed
20		failed

Table 8.

	$\alpha = \frac{3}{10}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$3.0E - 06$	3.906
10		failed
15		failed
20		failed

Table 7.

	$\alpha = \frac{1}{2}, \mu = \frac{1}{5}, \beta = \frac{1}{4}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$3.0E - 07$	1.375
10	$1.0E - 11$	1.672
15	$1.2E - 15$	2.328
20	$7.0E - 20$	3.266

Table 9.

$$L_0(t) = 1, \quad L_1(t) = 2t - 1. \quad (3.9)$$

Note that $L_n(0) = (-1)^n$ and $L_n(1) = 1$. The shifted fractional-order Legendre functions defined by introducing the change of variable $t = x^\alpha$ with $\alpha > 0$ on shifted Legendre polynomials, are denoted by $FL_i^\alpha(x)$ [10].

Hence $FL_i^\alpha(x)$ satisfy the following recurrence relation

$$FL_{n+1}^\alpha(x) = \frac{(2n+1)(2x^\alpha - 1)}{(n+1)} FL_n^\alpha(x) - \frac{n}{n+1} FL_{n-1}^\alpha(x), \quad n = 1, 2, 3, \dots,$$

$$FL_0^\alpha(x) = 1, \quad FL_1^\alpha(x) = 2x^\alpha - 1.$$

3.3 Collocation method

Consider the linear fractional differential equation:

$$\sum_{k=0}^n D^{\alpha_k} y(x) = g(x), \quad (3.10)$$

where $\alpha_k \in (k, k + 1]$, with initial conditions

$$y^{(i)}(0) = \beta_i, \quad i = 0, 1, \dots, n. \quad (3.11)$$

The unknown function $y(t)$ in problem (3.10), can be approximated by a truncated series of Legendre functions,

$$y_m(t) = \sum_{j=0}^m c_j FL_j^\alpha(t), \quad (3.12)$$

where c_j are unknowns. Here, the main purpose is to find c_j . In order to achieve this end, putting (3.12) in (3.10) and (3.11) we obtain:

$$\sum_{j=0}^m c_j \sum_{k=0}^n D^{\alpha_k} FL_j^\alpha(t) = g(t), \quad (3.13)$$

$$\sum_{j=0}^m c_j FL_j^{\alpha(i)}(0) = \beta_i, \quad i = 0, 1, \dots, n. \quad (3.14)$$

Relation (3.14) forms a system with $n + 1$ equations and $m + 1$ unknowns, to construct the remaining $m - n$ equations, we substitute Legendre-Guass points $\{t_i\}_{i=1}^{m-n}$ in (3.13), to obtain $m - n$ equations. So, reduces the obtaining to the solution of the system $AC = b$, where A, C and b are $A = \begin{bmatrix} A1 \\ A2 \end{bmatrix}$, $C = [c_0, c_1, \dots, c_m]^T$, $b = \begin{bmatrix} b1 \\ b2 \end{bmatrix}$ and matrices $A1_{(m-n) \times (m+1)}$ and $A2_{(n+1) \times (m+1)}$ are defined by

$$A1[i, j] = \sum_{k=0}^n D^{\alpha_k} FL_j^\alpha(t_i), \quad i = 1, 2, \dots, m - n,$$

$$j = 0, 1, \dots, m,$$

$$A2[i, j] = FL_j^{\alpha(i)}(0), \quad i = 0, 1, \dots, n,$$

$$j = 0, 1, \dots, m,$$

and vectors $b1_{(m-n) \times 1}, b2_{(n+1) \times 1}$ are defined by $b1[i] = g(t_i), \quad i = 1, 2, \dots, m - n,$
 $b2[i] = \beta_i, \quad i = 0, 1, \dots, n.$

	$\alpha = \frac{1}{4}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$	
L	$\ RESy\ _\infty$	Cpu Times
5	$3.0E - 06$	2.469
10	$1.2E - 09$	3.422
15	$5.0E - 13$	4.531
20	$1.6E - 16$	5.984

Table 10.

	$\alpha = \frac{4}{5}, \mu = \frac{1}{2}, \beta = \frac{1}{5}$	
L	$\ RESy\ _\infty$	CPU Times
5	$1.6E - 05$	6.766
10	$2.0E - 08$	12.250
15	$2.0E - 11$	18.953
20	$9.0E - 15$	23.984

Table 12.

	$\alpha = \frac{3}{10}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$	
L	$\ RESy\ _\infty$	CPU Times
5	$3.0E - 06$	2.515
10	$1.0E - 09$	3.516
15	$2.5E - 13$	4.828
20	$5.0E - 17$	7.110

Table 11.

3.4 The methodology

Consider the fractional logistic equation

$$D^\alpha y(x) = \mu y(x)(1 - y(x)), \quad (3.15)$$

where $\alpha \in (0, 1]$ [27], with initial condition

$$y(0) = \beta. \quad (3.16)$$

The nonlinear equation (3.15), can be written by

$$\begin{cases} D^\alpha y(x) - \mu y(x) = -\mu y^2(x), \\ y(0) = \beta. \end{cases} \quad (3.17)$$

Substituting the $y(x) = \sum_{k=0}^\infty y_k(x)$ in the non-linear fractional logistic equation, we have:

$$\begin{aligned} \sum_{k=0}^\infty D^\alpha y_k(x) - \sum_{k=0}^\infty \mu y_k(x) &= -\mu \left(\sum_{k=0}^\infty y_k(x) \right)^2 \\ &= -\sum_{k=0}^\infty A_k, \end{aligned}$$

where $A_0 = 0$ and

$$A_k = -\mu \left(\sum_{l=0}^{k-1} y_l \right)^2 + \mu \left(\sum_{l=0}^k y_l \right)^2.$$

The solution of problem (3.17), is

$$y(x) = \sum_{k=0}^\infty y_k(x)$$




Figure 1: Comparing the exact solution and approximate solution by NSIM and Adams method.

where $y_k(x)$ satisfies in

$$\begin{aligned} D^\alpha y_k(x) - \mu y_k(x) &= -A_k, \\ k &= 0, 1, \dots \end{aligned} \quad (3.18)$$

We solve the above linear equation using the spectral method. The function $y_k(x)$ can be approximated as

$$y_k(x) = \sum_{j=0}^\infty c_j^{(k)} FL_j^\alpha(x),$$

where the unknown coefficients $c_j^{(k)}$ are determined by using the collocation method. The residual function associated to the equation (3.18) is

$$\begin{aligned} RESy_k(x) &= D^\alpha y_k(x) - \mu y_k(x) + A_k, \\ k &= 0, 1, \dots \end{aligned}$$

By imposing the initial condition (3.16), we have

$$\sum_{j=0}^\infty c_j^{(k)} FL_j^\alpha(0) = \begin{cases} \beta, & k = 0, \\ 0, & k = 1, 2, \dots \end{cases}$$

For all k , the matrix form of the above system is:

$$MC^{(k)} = b^{(k)},$$




Figure 2: $\alpha = \frac{1}{2}, \mu = \frac{1}{5}, \beta = \frac{1}{4}$




Figure 4: $\alpha = \frac{3}{10}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$




Figure 3: $\alpha = \frac{1}{4}, \mu = \frac{1}{4}, \beta = \frac{1}{10}$




Figure 5: $\alpha = \frac{4}{5}, \mu = \frac{1}{2}, \beta = \frac{1}{5}$

$$M = [m_{ij}]_{(n+1)(n+1)},$$

$$C^{(k)} = [c_0^{(k)}, c_1^{(k)}, \dots, c_n^{(k)}]^t,$$

$$b^{(k)} = [b_0^{(k)}, b_1^{(k)}, \dots, b_n^{(k)}]^t.$$

Suppose that $\{x_i\}_{i=1}^n$ are zeros of Legendre polynomial of degree n, we have

$$m_{0j} = FL_j^\alpha(0),$$

$$j = 0, 1, 2, \dots, n,$$

$$m_{ij} = D^\alpha FL_j^\alpha(x_i) - \mu FL_j^\alpha(x_i),$$

$$j = 0, 1, 2, \dots, n, \quad i = 1, 2, \dots, n.$$

For $i = 1, 2, 3, \dots, n, k = 1, 2, 3, \dots$, we have

$$b_0^{(0)} = \beta, \quad b_i^{(0)} = 0,$$

and

$$b_0^{(k)} = 0, \quad b_i^{(k)} = -A_k(x_i).$$

The approximate solution of (3.15) with $L + 1$ terms is

$$y_{L,n} = \sum_{k=0}^L y_k(x), \tag{3.19}$$

where

$$y_k(x) = \sum_{j=0}^n c_j^{(k)} FL_j^\alpha(x). \tag{3.20}$$

4 Numerical study

Consider the following logistic initial value problem:

$$\begin{cases} D^\alpha y(x) = \mu y(x)(1 - y(x)), \\ y(0) = \beta. \end{cases} \tag{4.21}$$

We demonstrate the effectiveness of the proposed method (NSIM) by applying it on four values of α, β and μ for above problem. For each case, the maximum norm of the residual error of $y_{L,n}(x)$ is presented. Tables 1, 2, 3 and 4 shows the obtained numerical results of the (NSIM), tables 5, 6, 7 and 8 shows the obtained numerical results of the (IM) and tables 9, 10, 11 and 12 shows the obtained numerical results of the (ADM). The exact solution of (4.21) is $y(x) = \frac{e^{0.5x}}{1+e^{0.5x}}$ for $\alpha = 0.5$,

$\mu = 0.5$ and $\beta = 0.5$. The figure 1 shows the solutions obtained by (NSIM) and Adams method with $h = 0.001$ and exact solution.

All the computations associated with the method have been performed by a personal computer having the Intel Pentium 4, 2.8 GHz processor, 1GB RAM and using Maple 13 with 32 digits precision.

5 Estimation of the errors

The approximate solution of (4.21) is $y_{L,n}(x)$ and the exact solution is $y(x)$. Substituting $y_{L,n}(x)$ and $y(x)$ in (4.21), we obtain the following results.

$$\begin{cases} D^\alpha y(x) - \mu y(x)(1 - y(x)) = 0, \\ y(0) = \beta, \end{cases} \quad (5.22)$$

$$\begin{cases} D^\alpha y_{L,n}(x) - \mu y_{L,n}(x)(1 - y_{L,n}(x)) = R(x), \\ y_{L,n}(0) = \beta, \end{cases} \quad (5.23)$$

where $R(x)$ is the residual error. From (5.22) and (5.23) we obtain

$$\begin{cases} D^\alpha E_{L,n}(x) = \mu E_{L,n}(x)(1 + E_{L,n}(x) - 2y_{L,n}(x)) + R(x), \\ E_{L,n}(0) = 0, \end{cases} \quad (5.24)$$

where $E_{L,n} = y_{L,n}(x) - y(x)$ is error of solution. The solution of the (5.24) is an estimate of the error of $y_{L,n}(x)$. To have convergence we should have $|R(x)| \simeq |E_{L,n}(x)|$ and $\lim_{L,n \rightarrow \infty} |E_{L,n}(x)| = 0$.

We calculate $E_{L,n}(x)$ by Adams method for $h = 0.1$ and compare with residual error of NSIM for $L = 20$ in figures 2, 3, 4 and 5.

6 Conclusion

In this paper we proposed a new method to solve logistic equations of fractional order. This method was based on combination of iterative and spectral methods, which reduced nonlinear differential equations to systems of linear algebraic equations. The obtained approximate solutions have shown the effectiveness of our new method.

References

- [1] I. Podlubny, *Fractional Differential Equations*, Academic Press, San Diego, CA, 1999.
- [2] K. B. Oldham, J. Spanier, *The Fractional Calculus*, Academic Press, New York, London, 1974.
- [3] K. S. Miller, B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [4] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and applications of fractional differential equations*, Elsevier, Amsterdam, 2006.
- [5] R. Gorenflo, F. Mainardi, *Fractional calculus: integral and differential equations of fractional order in: A. Carpinteri, F. Mainardi (Eds.)*, Fractals and Fractional Calculus in Continuum Mechanics, Springer-Verlag, Wien/New York (1997) 223-276.
- [6] R. Gorenflo, A. Y. Luchko, *The initial value problem for some fractional differential equations with the Caputo derivative*, Fachbereich Mathematik und Informatik, Freie Universität Berlin, 1998.
- [7] F. Mainardi, *Fractional Calculus: some basic problems in continuum and statistical mechanics, in: A. Carpinteri, F. Mainardi (Eds.)*, Fractals and Fractional Calculus in Continuum Mechanics, Springer-Verlag, Wien/New York (1997) 291-348.
- [8] V. Daftardar-Gejji, H. Jafari, *Adomian decomposition: a tool for solving a system of fractional differential equations*, J. Math. Anal. Appl. 301 (2005) 508-518.
- [9] J. Shen, T. Tang, *High Order Numerical Methods and Algorithms*, Chinese Science Press, Beijing, 2005.
- [10] S. Kazem, S. Abbasbandy, Sunil Kumar, *Fractional-order Legendre functions for solving fractional-order differential equations*, Appl. Math. Modelling 37 (2013) 5498-5510.
- [11] K. T. Alligood, T. D. Sauer, J. A. Yorke, *An Introduction to Dynamical Systems*, Springer (1996).

- [12] M. Ausloos, *The Logistic map and the route to chaos: From the Beginnings to Modern Applications XVI*, 411 (2006).
- [13] R. L. Bagley, P. J. Torvik, *On the appearance of the fractional derivative in the behavior of real materials*, J. Appl. Mech. 51 (1984) 294-298.
- [14] J. M. Cushing, *An Introduction to Structured Population Dynamics*, Society for Industrial and Applied Mathematics (1998).
- [15] K. Diethelm, *An algorithm for the numerical solution of differential equations of fractional order*, Electron Trans. Numer. Anal. 5 (1997) 1-6.
- [16] A. M. A. El-Sayed, A. E. M. El-Mesiry, H. A. A. El-Saka, *On the fractional order Logistic equation*, Appl. Math. Letters 20 (2007) 817-823.
- [17] J. H. He, *Variational iteration method-a kind of non-linear analytical technique: some examples*, International Journal of Non-Linear Mechanics 34 (1999) 699-708.
- [18] M. M. Khader, *On the numerical solutions for the fractional diffusion equation*, Communications in Nonlinear Science and Numerical Simulation 16 (2011) 2535-2542.
- [19] M. M. Khader, *Introducing an efficient modification of the variational iteration method by using Chebyshev polynomials*, Application and Applied Mathematics: An International Journal 7 (2012) 283-299.
- [20] M. M. Khader, *Introducing an efficient modification of the homotopy perturbation method by using Chebyshev polynomials*, Arab Journal of Mathematical Sciences 18 (2012) 61-71.
- [21] M. M. Khader, N. H. Sweilam, A. M. S. Mahdy, *An efficient numerical method for solving the fractional diffusion equation*, Journal of Applied Mathematics and Bioinformatics 1 (2011) 1-12.
- [22] M. M. Khader, A. S. Hendy, *The approximate and exact solutions of the fractional-order delay differential equations using Legendre pseudospectral method*, International Journal of Pure and Applied Mathematics 74 (2012) 287-297.
- [23] H. Pastijn, *Chaotic Growth with the Logistic Model of P.-F. Verhulst, Understanding Complex Systems, (2006)*, The Logistic Map and the Route to Chaos, Pages 3-11.
- [24] N. H. Sweilam, M. M. Khader, R. F. Al-Bar, *Numerical studies for a multiorde differential equation*, Physics Letters A 371 (2007) 26-33.
- [25] N. H. Sweilam, M. M. Khader, *On the convergence of VIM for nonlinear coupled system of partial differential equations*, Int. J. of Computer Maths. 87 (2010) 1120-1130.
- [26] N. H. Sweilam, M. M. Khader, A. M. S. Mahdy, *Numerical studies for fractional-order Logistic differential equation with two different delays*, Accepted in Journal of Applied Mathematics, to appear in 2012.
- [27] S. Bhalekar, V. Daftardar-Gejji, *Solving Fractional-Order Logistic Equation Using a New Iterative Method*, International Journal of Differential Equations(Hindawi), (2012).
- [28] P. F. Verhulst, *Notice sur la loi que la population suit dans son accroissement*, Correspondence Math. Phys. 10 (1838) 113-121.
- [29] L. R. Devaney, *An introduction to chaotic dynamical system*, Benjamin 1985.
- [30] S. Bhalekar, V. Daftardar-Gejji, *Solving a System of Nonlinear Functional Equations Using Revised New Iterative Method*, International Journal of Computational and Mathematical Sciences 6 (2012).
- [31] H. Jafari, S. Seifi, *An Iterative Method for Solving a System of Nonlinear Algebraic Equations*, Journal of Applied Mathematics, Islamic Azad University of Lahijan 5 (2008).
- [32] S. Bhalekar, V. Daftardar-Gejji, *Convergence of the New Iterative Method*, International Journal of Differential Equations Volume 2011, Article ID 989065.
- [33] H. Jafari, M. Ahmadi, and S. Sadeghi, *Solving Singular Boundary Value Problems Using Daftardar-Jafari Method*, Applications

and Applied Mathematics: An International Journal (AAM) 7 (2012) 357-364.

- [34] M. Aslam. Noor, K. Inayat. Noor, E. Al-Said, M. Waseem, *Some New Iterative Methods for Nonlinear Equations*, Mathematical Problems in Engineering Volume 2010, Article ID 198943.
- [35] V. Daftardar-Gejji, H. Jafari, *An iterative method for solving nonlinear functional equations*, Journal of Mathematical Analysis and Applications 316 (2006) 753-763.
- [36] K. Diethelm, *An algorithm for the numerical solution of differential equations of fractional order*, Elec. Transact. Numer. Anal. 5 (1997) 1-6.



Ahmad Shoja is currently a Ph.D. student at Science and Research Branch, Islamic Azad University, Tehran, Iran. His research interests include the numerical solving of fractional differential equation with Spectral-Iterative methods.



Esmail Babolian is professor of applied mathematics and Faculty of Mathematical sciences and computer, Kharazmy University, Tehran, Iran. Interested in numerical solution of functional Equations, numerical linear algebra and mathematical education.



Alireza Vahidi is associated professor of applied mathematics. Yadegare-Emam Branch, Islamic Azad University, Tehran, Iran. Interested in numerical solution of integral and differential equations.